

A HIERARCHICAL MODEL OF QUANTUM ANHARMONIC OSCILLATORS: CRITICAL POINT CONVERGENCE

S. ALBEVERIO, YU. KONDRATIEV, A. KOZAK, AND YU. KOZITSKY

to appear in Communications in Mathematical Physics

ABSTRACT. A hierarchical model of interacting quantum particles performing anharmonic oscillations is studied in the Euclidean approach, in which the local Gibbs states are constructed as measures on infinite dimensional spaces. The local states restricted to the subalgebra generated by fluctuations of displacements of particles are in the center of the study. They are described by means of the corresponding temperature Green (Matsubara) functions. The result of the paper is a theorem, which describes the critical point convergence of such Matsubara functions in the thermodynamic limit.

CONTENTS

1.	Introduction	1
2.	Setup and the Theorem	5
3.	Euclidean Representation	8
4.	Proof of the Theorem	13
5.	Proof of Lemma 4.1	15
	References	22

1. INTRODUCTION

Let \mathbb{L} be a countable set (lattice). With each $l \in \mathbb{L}$ we associate a quantum mechanical particle with one degree of freedom described by the momentum \mathfrak{p}_l and displacement \mathfrak{q}_l operators. The system of such particles which we consider in this article is described by the heuristic Hamiltonian

$$(1.1) \quad H = -\frac{1}{2} \sum_{l,l'} J_{ll'} \mathfrak{q}_l \mathfrak{q}_{l'} + \sum_l \left[\frac{1}{2m} \mathfrak{p}_l^2 + \frac{a}{2} \mathfrak{q}_l^2 + b \mathfrak{q}_l^4 \right].$$

Here $b > 0$, $a \in \mathbb{R}$ and the sums run through the lattice \mathbb{L} . The operators \mathfrak{p}_l and \mathfrak{q}_l satisfy the relation

$$(1.2) \quad [\mathfrak{p}_l, \mathfrak{q}_l] = \mathfrak{p}_l \mathfrak{q}_l - \mathfrak{q}_l \mathfrak{p}_l = 1/i,$$

1991 *Mathematics Subject Classification.* 82B27; 82B10.

Key words and phrases. Euclidean Gibbs states; abnormal fluctuations; criticality; quantum crystal.

Supported by Deutsche Forschungsgemeinschaft through the German-Polish project 436 POL 113/98/0-1 "Probability measures"; Agnieszka Kozak and Yuri Kozitsky were also supported by Komitet Badań Naukowych through the Grant 2P03A 02025.

and $\mathbf{m} = \mathbf{m}^{\text{phys}}/\hbar^2$ is the reduced mass of the particle. Models like (1.1) have been studied for many years, see e.g., [23, 29]. They (and their simplified versions) are used as a base of models describing strong electron-electron correlations caused by the interaction of electrons with vibrating ions [14, 30].

Let $\mathcal{L} = \{\Lambda_n\}_{n \in \mathbb{N}_0}$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ be a sequence of finite subsets of \mathbb{L} , which is ordered by inclusion and exhausts \mathbb{L} . For every Λ_n , let H_{Λ_n} be a local Hamiltonian, corresponding to (1.1). In a standard way the Hamiltonians H_{Λ_n} determine local Gibbs states $\varrho_{\beta, \Lambda_n}$. A phase transition in the model (1.1) is connected with macroscopic displacements of particles from their equilibrium positions $\mathbf{q}_l = 0$, $l \in \mathbb{L}$. To describe this phenomenon, one considers fluctuation operators

$$(1.3) \quad Q_{\Lambda_n}^{(\alpha)} \stackrel{\text{def}}{=} \frac{1}{|\Lambda_n|^{(1+\alpha)/2}} \sum_{l \in \Lambda_n} \mathbf{q}_l, \quad \alpha \geq 0,$$

and Matsubara functions

$$(1.4) \quad \Gamma_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) \stackrel{\text{def}}{=} \varrho_{\beta, \Lambda_n} \left\{ Q_{\Lambda_n}^{(\alpha)} \exp(-(\tau_2 - \tau_1)H_{\Lambda_n}) \cdots \right. \\ \left. \times \exp(-(\tau_{2k} - \tau_{2k-1})H_{\Lambda_n}) Q_{\Lambda_n}^{(\alpha)} \exp((\tau_{2k} - \tau_1)H_{\Lambda_n}) \right\}, \quad k \in \mathbb{N},$$

with the arguments satisfying the condition $0 \leq \tau_1 \leq \dots \leq \tau_{2k} \leq \beta$. In our model the interaction potential is taken to be

$$(1.5) \quad J_{ll'} = J[d(l, l') + 1]^{-1-\delta}, \quad J, \delta > 0,$$

where $d(l, l')$ is a metric on \mathbb{L} , determined by means of a hierarchical structure. The latter is a family of finite subsets of \mathbb{L} , each of which belongs to a certain hierarchy level $n \in \mathbb{N}_0$. This fact predetermines also our choice of the sequence \mathcal{L} – the subsets Λ_n are to be the elements of the hierarchical structure, that is typical for proving scaling limits in hierarchical models (see e.g., [12]). We prove (Theorem 2.2) that, for any $\delta \in (0, 1/2)$, the parameters $a \in \mathbb{R}$, $b > 0$ and $\mathbf{m} > 0$ can be chosen in such a way that there will exist $\beta_* > 0$ with the following properties:

- (a) if $\beta = \beta_*$, for all $k \in \mathbb{N}$, the functions (1.4) converge

$$(1.6) \quad \Gamma_{2k}^{\delta, \beta_*, \Lambda_n}(\tau_1, \dots, \tau_{2k}) \longrightarrow \frac{(2k)!}{k! 2^k \beta_*^k} \left(\frac{J_*}{J} \right)^k, \quad n \rightarrow +\infty$$

uniformly with respect to their arguments; here $J_* > 0$ is a constant determined by the hierarchical structure;

- (b) if $\beta < \beta_*$, for all $\alpha > 0$ and $k \in \mathbb{N}$, the functions $\Gamma_{2k}^{\alpha, \beta, \Lambda_n}$ converge to zero in the same sense.

The convergence of the functions (1.4) like in (1.6) but with $\alpha = 1$ would correspond to the appearance of a long-range order, which destroys the Z_2 -symmetry. Thus, claim (a) describes a critical point where the fluctuations are abnormal (since $\alpha = \delta > 0$) but not strong enough to destroy the mentioned symmetry. Such fluctuations are classical (non-quantum), which follows from the fact that the limits (1.6) are independent of τ .

Due to the hierarchical structure the model (1.1) is self-similar. In translation invariant lattice models self-similarity appears at their critical points [27, 28]. This, among others, is the reason why the critical point properties of hierarchical models

of classical statistical mechanics attract attention during the last three decades. An expository review of the results in this domain is given in [12].

In the model (1.1) the oscillations are described by unbounded operators¹. The same model was studied in our previous works [2, 3, 4]. In [2] a preliminary study of the model was performed. A theorem describing the critical point convergence was announced in [3]. In [4] we have shown that the critical point of the model (1.1) can be suppressed by strong quantum effects, which take place, in particular, when the mass m is less than a certain bound m_* ². In the present paper we give a complete proof of the critical point convergence, which appears for sufficiently large values of the mass (see the discussion at the very end of this introduction). It should be pointed out that, to the best of our knowledge, our result is the first example of a theorem, which describes the convergence at the critical point of a nontrivial quantum model, published by this time.

Let us outline the main aspects of the proof. By symmetry, the functions (1.4) are extended to \mathcal{I}_β^{2k} , where $\mathcal{I}_\beta \stackrel{\text{def}}{=} [0, \beta]$. Then for $x \in L^2(\mathcal{I}_\beta)$, one sets

$$(1.7) \quad \varphi_n^{(\alpha)}(x) = 1 + \sum_{k=1}^{\infty} \frac{1}{(2k)!} \int_{\mathcal{I}_\beta^{2k}} \Gamma_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) x(\tau_1) \cdots x(\tau_{2k}) d\tau_1 \cdots d\tau_{2k},$$

and

$$(1.8) \quad \log \varphi_n^{(\alpha)}(x) = \sum_{k=1}^{\infty} \frac{1}{(2k)!} \int_{\mathcal{I}_\beta^{2k}} U_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) x(\tau_1) \cdots x(\tau_{2k}) d\tau_1 \cdots d\tau_{2k},$$

which uniquely determines the Ursell functions $U_{2k}^{\alpha, \beta, \Lambda_n}$. In terms of these functions our result may be formulated as follows:

$$(1.9) \quad \begin{aligned} U_2^{\delta, \beta_*, \Lambda_n}(\tau_1, \tau_2) &\longrightarrow \beta_*^{-1}, \\ \forall k > 1 : U_{2k}^{\delta, \beta_*, \Lambda_n}(\tau_1, \dots, \tau_{2k}) &\longrightarrow 0, \\ \forall k \in \mathbb{N}, \beta < \beta_*, \alpha > 0 : U_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) &\longrightarrow 0, \end{aligned}$$

which holds uniformly with respect to the arguments $\tau_j \in \mathcal{I}_\beta$, $j = 1, \dots, 2k$ as $n \rightarrow +\infty$. Here we have set $J = J_*$, that can always be done by choosing an appropriate scale of β . We prove (1.9) in the framework of the Euclidean approach in quantum statistical mechanics based on the representation of the functions (1.4) in the form of functional integrals. This approach was initiated in [1, 15], its detailed description and an extended related bibliography may be found in [6]. In separate publications we are going to exploit our result, in particular, to construct self-similar Gibbs states (in the spirit of [11, 12] where it was done for classical hierarchical models).

The functions $\Gamma_{2k}^{\alpha, \beta, \Lambda_n}$, $U_{2k}^{\alpha, \beta, \Lambda_n}$, $k \in \mathbb{N}$ are continuous on \mathcal{I}_β^{2k} , see [6]. In view of our choice of the potential energy in (1.1), the Ursell functions satisfy the sign rule

$$(1.10) \quad (-1)^{k-1} U_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) \geq 0,$$

for all $k \in \mathbb{N}$ and $(\tau_1, \dots, \tau_{2k}) \in \mathcal{I}_\beta^{2k}$. We prove that the families $\{\Gamma_{2k}^{\alpha, \beta, \Lambda_n}\}_{n \in \mathbb{N}_0}$, $\{U_{2k}^{\alpha, \beta, \Lambda_n}\}_{n \in \mathbb{N}_0}$, $k \in \mathbb{N}$ are equicontinuous; hence, the convergence (1.9) can be

¹Certain aspects of critical point behaviour of quantum hierarchical models with bounded (spin) operators were studied in [20].

²Physical aspects of such quantum effects were analyzed in [7].

proven by showing the convergence of $U_2^{\delta, \beta_*, \Lambda_n}$, $U_2^{\alpha, \beta, \Lambda_n}$, as in (1.9), and

$$(1.11) \quad \mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n} \stackrel{\text{def}}{=} \int_{\mathcal{I}_\beta^{2k}} U_{2k}^{\alpha, \beta, \Lambda_n}(\tau_1, \dots, \tau_{2k}) d\tau_1 \cdots d\tau_{2k} \longrightarrow 0,$$

which has to hold for all $k \geq 2$, $\beta \leq \beta_*$, and for $\alpha = \delta$ if $\beta = \beta_*$, and $\alpha > 0$ if $\beta < \beta_*$. Another fact which we employ here is also a consequence of the choice of the potential energy in (1.1). By a version of the Lee-Yang theorem, the function f_n of a single complex variable defined in the vicinity of $z = 0$ by the series

$$(1.12) \quad \log f_n(z) = \sum_{k=1}^{\infty} \frac{1}{(2k)!} \mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n} z^{2k},$$

can be extended to an even entire function of order less than two possessing imaginary zeros only. This implies

$$(1.13) \quad f_n(z) = \prod_{j=1}^{\infty} (1 + c_j^{(n)} z^2), \quad c_1^{(n)} \geq c_2^{(n)} \geq \cdots > 0, \quad \sum_{j=1}^{\infty} c_j^{(n)} < \infty,$$

yielding for the numbers (1.11) the following representation

$$(1.14) \quad \mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n} = 2(2k-1)!(-1)^{k-1} \sum_{j=1}^{\infty} [c_j^{(n)}]^k, \quad k \in \mathbb{N},$$

by which,

$$(1.15) \quad |\mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n}| \leq 2(2k-1)! [c_1^{(n)}]^{k-2} \sum_{j=1}^{\infty} [c_j^{(n)}]^2, \quad k \geq 2,$$

$$|\mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n}| \leq (2k-1)! [c_1^{(n)}]^{k-1} \mathcal{U}_2^{\alpha, \beta, \Lambda_n} \quad k \in \mathbb{N},$$

and hence

$$(1.16) \quad |\mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n}| \leq (2k-1)!(2^{1-k}/3) [\mathcal{U}_2^{\alpha, \beta, \Lambda_n}]^{k-2} |\mathcal{U}_4^{\alpha, \beta, \Lambda_n}|, \quad k \geq 2,$$

$$|\mathcal{U}_{2k}^{\alpha, \beta, \Lambda_n}| \leq (2k-1)! 2^{1-k} [\mathcal{U}_2^{\alpha, \beta, \Lambda_n}]^k, \quad k \in \mathbb{N}.$$

Then the convergence (1.9) follows from the corresponding convergence of $U_2^{\alpha, \beta, \Lambda_n}$ and from the fact

$$(1.17) \quad \mathcal{U}_4^{\delta, \beta_*, \Lambda_n} \longrightarrow 0.$$

The above arguments allow us to prove the convergence of an infinite number of sequences of functions by controlling just two sequences of numbers – $\{\hat{u}_n\}_{n \in \mathbb{N}_0}$ and $\{\mathcal{U}_4^{\delta, \beta_*, \Lambda_n}\}_{n \in \mathbb{N}_0}$, where $\hat{u}_n = \beta^{-1} \mathcal{U}_2^{\delta, \beta, \Lambda_n}$. The sign rule (1.10) and the representation (1.14) are proven in Lemmas 3.3 and 3.4 below by means of the lattice approximation technique [6]. Here the functions $\Gamma_{2k}^{\alpha, \beta, \Lambda_n}$, $k \in \mathbb{N}$, are obtained as limits of moments of Gibbs measures of classical ferromagnetic ϕ^4 -models. This allows us to employ the corresponding properties of the ϕ^4 -models proven in [24] (the sign rule), [2] (a correlation inequality) and [19] (the Lee-Yang theorem). Then to controlling the sequences $\{\hat{u}_n\}_{n \in \mathbb{N}_0}$ and $\{\mathcal{U}_4^{\delta, \beta_*, \Lambda_n}\}_{n \in \mathbb{N}_0}$ we apply a version of the inductive method developed in [17, 18]. The central role here is played by Lemma 4.1. It establishes the existence of $\beta_* > 0$ such that, for $\beta = \beta_*$ (respectively, for $\beta < \beta_*$), the sequence $\{\hat{u}_n\}_{n \in \mathbb{N}_0}$ converges to one (respectively, to zero as

$|\Lambda_n|^{-\delta}$). The sequence $\{\mathcal{U}_4^{\delta,\beta,\Lambda_n}\}_{n \in \mathbb{N}_0}$ converges to zero in both cases. The latter fact is proven by constructing a converging to zero sequence of positive numbers $\{X_n\}_{n \in \mathbb{N}_0}$, such that $\beta^{-2}|\mathcal{U}_4^{\delta,\beta,\Lambda_n}| \leq X_n$ for all $n \in \mathbb{N}_0$ and $\beta \leq \beta_*$. The proof of Lemma 4.1 is based on recurrent estimates (Lemma 5.1) yielding upper and lower bounds for \hat{u}_n and X_n in terms of certain functions of \hat{u}_{n-1} and X_{n-1} . The analysis of these estimates shows that the simultaneous convergence $\hat{u}_n \rightarrow 1$ and $X_n \rightarrow 0$ can be guaranteed if these sequences are confined to the intervals $\hat{u}_n \in (1, \bar{v})$ and $X_n \in (0, \bar{w})$, where the parameters $\bar{v} > 1$ and $\bar{w} > 0$ depend on δ and on the details of the hierarchical structure only and can be computed explicitly. Lemma 5.1 is proven by comparing solutions of certain differential equations, similarly as in [17, 18]. Lemma 5.4 establishes the existence of $\beta_n^\pm > 0$, $\beta_n^- < \beta_n^+$ if β_{n-1}^\pm do exist. These numbers are defined as follows: $\hat{u}_n = \bar{v}$ for $\beta = \beta_n^+$, and $\hat{u}_n < \bar{v}$ for $\beta < \beta_n^+$; $\hat{u}_n = 1$ for $\beta = \beta_n^-$, and $\hat{u}_n < 1$ for $\beta < \beta_n^-$. The proof of Lemma 5.4 is carried out by means of the estimates obtained in Lemma 5.1. In Lemma 5.3 we prove that the parameters \mathfrak{m} , a and b can be chosen in such a way that β_0^\pm do exist. In Lemma 5.5 we prove the existence of β_* , such that $\forall n \in \mathbb{N}_0 : \hat{u}_n \in (1, \bar{v})$ for $\beta = \beta_*$, and $\hat{u}_n \rightarrow 0$ as $|\Lambda_n|^{-\delta}$ for $\beta < \beta_*$. The proof is based on the estimates obtained in Lemma 5.1. In Lemma 3.5 we prove that all \hat{u}_n , $n \in \mathbb{N}_0$ are continuous functions of β and describe certain useful properties of the Ursell functions $U_2^{\alpha_*,\beta,\Lambda_n}(\tau, \tau')$, $n \in \mathbb{N}_0$, implying e.g., the mentioned equicontinuity.

The proof of Lemma 5.3 is based on the estimates of \hat{u}_0 and X_0 obtained in Lemma 5.2. In particular, we prove that

$$\frac{\mathfrak{m}\gamma^2}{36} \left[1 - \exp\left(-\frac{3\beta}{\mathfrak{m}\gamma}\right) \right] \leq \hat{u}_0 \leq \frac{\beta\gamma}{8} \left[1 + \sqrt{1 + \frac{16}{\beta\gamma}} \right],$$

where $a < 0$, $\gamma = |a|/b$. Then for $\mathfrak{m}\gamma^2 > 36\bar{v}$, one gets $\hat{u}_0 > \bar{v}$ for sufficiently large β . On the other hand, $\hat{u}_0 \rightarrow 0$ as $\beta \rightarrow 0$. Since \hat{u}_0 depends on β continuously, this yields the existence of β_0^\pm . Furthermore, for fixed γ and β , we show that $X_0 \leq bC$ with a certain fixed $C > 0$. This was used to provide $X_0 < \bar{w}$, and hence $X_n < \bar{w}$, $n \in \mathbb{N}$, for sufficiently small $b > 0$. Another upper bound of \hat{u}_0 was obtained in [4]. It is well-known that the one particle Hamiltonian which stands in the square brackets in (1.1) has a pure point non-degenerate spectrum. Let E_n , $n \in \mathbb{N}_0$ be its eigenvalues and $\Delta = \min_{n \in \mathbb{N}}(E_n - E_{n-1})$. In [4] we proved that if $\mathfrak{m}\Delta^2 > 1$, then $\hat{u}_0 < 1$ and hence $\hat{u}_n \rightarrow 0$ for all β . In what follows, the critical point of the model exists if $a < 0$ and the parameters $\mathfrak{m}(|a|/b)^2$, $1/b$ are big enough; such a point does not exist if ‘the quantum rigidity’ $\mathfrak{m}\Delta^2$ (see [7]) is greater than 1. By Lemma 1.1 of [4], $\mathfrak{m}\Delta^2 \sim \mathfrak{m}^{-1/3}C$, $C > 0$ as $\mathfrak{m} \rightarrow 0$, which means that small values of the mass prevent the system from criticality.

2. SETUP AND THE THEOREM

Like in [3, 4] we consider the hierarchical model defined on $\mathbb{L} = \mathbb{N}_0$. Given $\varkappa \in \mathbb{N} \setminus \{1\}$, we set

$$(2.1) \quad \Lambda_{n,s} = \{l \in \mathbb{N}_0 \mid \varkappa^n s \leq l \leq \varkappa^n(s+1) - 1\}, \quad s, n \in \mathbb{N}_0.$$

Then for $n \in \mathbb{N}$, one has

$$(2.2) \quad \Lambda_{n,s} = \bigcup_{l \in \Lambda_{k,s}} \Lambda_{n-k,l}, \quad k = 1, 2, \dots, n.$$

The collection of families $\{\Lambda_{n,s}\}_{s \in \mathbb{N}_0}$, $n \in \mathbb{N}_0$ is called a *hierarchical structure* on \mathbb{L} . Given $l, l' \in \mathbb{L}$, we set

$$(2.3) \quad n(l, l') = \min\{n \mid \exists \Lambda_{n,s} : l, l' \in \Lambda_{n,s}\}, \quad d(l, l') = \varkappa^{n(l, l')} - 1.$$

The function $d : \mathbb{L} \times \mathbb{L} \rightarrow [0, +\infty)$ has the following property: any triple $\{l_1, l_2, l_3\} \subset \mathbb{L}$ contains two elements, say l_1, l_2 , such that $d(l_1, l_3) = d(l_2, l_3)$. Thus, $d(l, l')$ is a metric on \mathbb{L} . The interaction potential in our model has the form of (1.5) with the above metric $d(l, l')$. It is invariant under the transformations of \mathbb{L} which leave $d(l, l')$ unchanged. In view of this fact, it is convenient to choose the sequences \mathcal{L} which determines the infinite-volume limit to be consisting of the sets (2.2) only.

A standard choice is the sequence of $\Lambda_{n,0} \stackrel{\text{def}}{=} \Lambda_n$, $n \in \mathbb{N}_0$.

The Hamiltonian (1.1) may be rewritten in the form

$$(2.4) \quad H = -\frac{\theta}{2} \sum_{n=0}^{\infty} \varkappa^{-n(1+\delta)} \sum_{l \in \mathbb{L}} \left(\sum_{l' \in \Lambda_{n,l}} q_{l'} \right)^2 + \sum_{l \in \mathbb{L}} \left[\frac{1}{2m} p_l^2 + \frac{a}{2} q_l^2 + b q_l^4 \right],$$

where $\theta = J(1 - \varkappa^{-(1+\delta)}) > 0$. The local Hamiltonians indexed by $\Lambda_{n,l}$ are obtained from the above one by the corresponding truncation of the sums. For our purposes, it is convenient to write them recursively

$$(2.5) \quad H_{\Lambda_{n,l}} \stackrel{\text{def}}{=} H_{n,l} = -\frac{\theta}{2} \varkappa^{-n(1+\delta)} \left(\sum_{s \in \Lambda_{n,l}} q_s \right)^2 + \sum_{s \in \Lambda_{1,l}} H_{n-1,s},$$

where the one particle Hamiltonian is

$$(2.6) \quad H_{0,l} = \frac{1}{2m} p_l^2 + \frac{a}{2} q_l^2 + b q_l^4.$$

The canonical pair p_l, q_l , as well as the Hamiltonian $H_{0,l}$, are defined in the complex Hilbert space $\mathcal{H}_l = L^2(\mathbb{R})$ as unbounded operators, which are essentially self-adjoint on the dense domain $C_0^\infty(\mathbb{R})$. The Hamiltonian $H_{n,l}$, $n \in \mathbb{N}$ is defined similarly but in the space $\mathcal{H}_{n,l} = L^2(\mathbb{R}^{|\Lambda_{n,l}|})$.

The local Gibbs state in $\Lambda_{n,l}$ at a given temperature $\beta^{-1} > 0$ is defined on $\mathfrak{C}_{n,l}$ – the C^* -algebra of bounded operators on $\mathcal{H}_{n,l}$, as follows

$$(2.7) \quad \varrho_{\beta, \Lambda_{n,l}}(A) = \frac{\text{trace}(A \exp(-\beta H_{n,l}))}{\text{trace} \exp(-\beta H_{n,l})}, \quad A \in \mathfrak{C}_{n,l}.$$

In a standard way, it may be extended to unbounded operators such as $q_{l'}, l' \in \Lambda_{n,l}$. The dynamics in $\Lambda_{n,l}$ is described by the time automorphisms of $\mathfrak{C}_{n,l}$

$$(2.8) \quad \alpha_{n,l}^t(A) = \exp(itH_{n,l}) A \exp(-itH_{n,l}), \quad t \in \mathbb{R}.$$

For a measurable function $A : \mathbb{R}^{|\Lambda_{n,l}|} \rightarrow \mathbb{C}$, the multiplication operator A acts on $\psi \in \mathcal{H}_{n,l}$ as

$$(A\psi)(x) = A(x)\psi(x), \quad x \in \mathbb{R}^{|\Lambda_{n,l}|}.$$

It appears that the linear span of the operators

$$\alpha_{n,l}^{t_1}(A_1) \cdots \alpha_{n,l}^{t_k}(A_k), \quad k \in \mathbb{N}, \quad t_1, \dots, t_k \in \mathbb{R},$$

with all possible choices of k, t_1, \dots, t_k and multiplication operators $A_1, \dots, A_k \in \mathfrak{C}_{n,l}$ is dense in the algebra $\mathfrak{C}_{n,l}$ in the σ -weak topology, in which the state (2.7) is

continuous. Thus, this state is fully determined by temporal Green functions

$$(2.9) \quad G_{A_1, \dots, A_k}^{n,l}(t_1, \dots, t_k) = \varrho_{\beta, \Lambda_{n,l}}(\mathbf{a}_{n,l}^{t_1}(A_1) \dots \mathbf{a}_{n,l}^{t_k}(A_k)),$$

corresponding to all possible multiplication operators $A_1, \dots, A_k \in \mathfrak{C}_{n,l}$. Set

$$(2.10) \quad \mathcal{D}_k^\beta = \{(t_1, \dots, t_k) \in \mathbb{C}^k \mid 0 < \text{Im}(t_1) < \dots < \text{Im}(t_k) < \beta\}.$$

As was proven in Lemma 2.1 in [6], every Green function (2.9) may be extended to a holomorphic function on \mathcal{D}_k^β . This extension is continuous on the closure of \mathcal{D}_k^β and may be uniquely determined by its values on the set

$$(2.11) \quad \mathcal{D}_k^\beta(0) = \{(t_1, \dots, t_k) \in \mathcal{D}_k^\beta \mid \text{Re}(t_j) = 0, \quad j = 1, \dots, k\}.$$

The restriction of $G_{A_1, \dots, A_k}^{n,l}$ to $\mathcal{D}_k^\beta(0)$, i.e., the function

$$(2.12) \quad \Gamma_{A_1, \dots, A_k}^{n,l}(\tau_1, \dots, \tau_k) = G_{A_1, \dots, A_k}^{n,l}(i\tau_1, \dots, i\tau_k),$$

is the Matsubara function corresponding to the operators A_1, \dots, A_k . By (2.7) - (2.9), it may be written

$$(2.13) \quad \begin{aligned} \Gamma_{A_1, \dots, A_k}^{n,l}(\tau_1, \dots, \tau_k) &= \frac{1}{Z_{n,l}} \text{trace} \{A_1 \exp(-(\tau_2 - \tau_1)H_{n,l}) \\ &\quad \times A_2 \exp(-(\tau_3 - \tau_2)H_{n,l}) \dots A_k \exp(-(\beta - \tau_k + \tau_1)H_{n,l})\}; \\ Z_{n,l} &\stackrel{\text{def}}{=} \text{trace} \{\exp(-\beta H_{n,l})\}. \end{aligned}$$

This representation immediately yields the ‘KMS-periodicity’

$$(2.14) \quad \Gamma_{A_1, \dots, A_k}^{n,l}(\tau_1 + \vartheta, \dots, \tau_k + \vartheta) = \Gamma_{A_1, \dots, A_k}^{n,l}(\tau_1, \dots, \tau_k),$$

for every $\vartheta \in \mathcal{I}_\beta \stackrel{\text{def}}{=} [0, \beta]$, where addition is of modulo β .

As was mentioned in the introduction, the phase transition in the model is connected with the appearance of macroscopic displacements of particles from their equilibrium positions $\mathbf{q}_l = 0$, which occur when the fluctuations of such displacements become large. To describe them, we set (c.f., (1.3))

$$(2.15) \quad Q_{n,l}^\lambda = \frac{\lambda_n}{\sqrt{|\Lambda_{n,l}|}} \sum_{l' \in \Lambda_{n,l}} \mathbf{q}_{l'} = \frac{\lambda_n}{\varkappa^{n/2}} \sum_{l' \in \Lambda_{n,l}} \mathbf{q}_{l'},$$

where $\{\lambda_n\}_{n \in \mathbb{N}_0}$ is a sequence of positive numbers. The operators $Q_{n,l}^\lambda$ are unbounded, nevertheless, the corresponding Matsubara functions still possess almost all of those ‘nice’ properties which they have in the case of bounded operators. The next statement follows directly from Corollary 4.1 and Theorem 4.2 of [6].

Proposition 2.1. *For every $n \in \mathbb{N}_0$ and $k \in \mathbb{N}$, the functions $\Gamma_{Q_{n,l}^\lambda, \dots, Q_{n,l}^\lambda}^{n,l}$ are continuous on \mathcal{I}_β^{2k} , they can be analytically continued to the domains \mathcal{D}_{2k}^β .*

The convergence of the sequence $\{\Gamma_{Q_{n,j}^\lambda, \dots, Q_{n,j}^\lambda}^{n,j}\}_{n \in \mathbb{N}_0}$ with $\lambda_n = \varkappa^{-n/2}$, to a nonzero limit would mean the appearance of the long range order caused by macroscopic displacements of particles. The convergence with a slower decaying sequence $\{\lambda_n\}$ corresponds to the presence of a critical point.

Our model is described by the following parameters: $\delta > 0$, which determines the decay of the potential $J_{l'l'}$, see (1.5); $\theta \geq 0$, which determines its strength; the

mass \mathbf{m} and the parameters of the potential energy a and b , see e.g., (1.1). Since the choice of θ determines only the scale of β , we may set

$$(2.16) \quad \theta = \varkappa^\delta - 1,$$

which corresponds to the choice (see (1.6))

$$J = J_* \stackrel{\text{def}}{=} \frac{\varkappa^\delta - 1}{1 - \varkappa^{-1-\delta}}.$$

To simplify notations we write the operator (2.15) with $\lambda_n = \varkappa^{-n\delta/2}$ as $Q_{n,l}$ and

$$(2.17) \quad \Gamma_{Q_{n,l}, \dots, Q_{n,l}}^{n,l}(\tau_1, \dots, \tau_{2k}) = \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}).$$

Theorem 2.2. *For the model (1.1) with $\delta \in (0, 1/2)$, one can choose the parameters a , b and \mathbf{m} in such a way that there will exist $\beta_* > 0$, dependent on a , b , \mathbf{m} , with the following properties: (a) if $\beta = \beta_*$, then for all $k \in \mathbb{N}$, the convergence*

$$(2.18) \quad \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) \longrightarrow \frac{(2k)!}{k! 2^k \beta_*^k},$$

holds uniformly on $(\tau_1, \dots, \tau_{2k}) \in \mathcal{I}_\beta^{2k}$; (b) if $\beta < \beta_$, the functions $\Gamma_{2k}^{\alpha, \beta, \Lambda_{n,l}}$, $k \in \mathbb{N}$ defined by (1.4) converge to zero in the same sense for all $\alpha > 0$.*

3. EUCLIDEAN REPRESENTATION

In the Euclidean approach [6] the functions (2.12) corresponding to the multiplication operators A_1, \dots, A_{2k} , are written as follows

$$(3.1) \quad \Gamma_{A_1, \dots, A_{2k}}^{n,l}(\tau_1, \dots, \tau_{2k}) = \int_{\Omega_{n,l}} A_1(\omega_{n,l}(\tau_1)) \dots A_{2k}(\omega_{n,l}(\tau_{2k})) \nu_{n,l}(d\omega_{n,l}),$$

where $\Omega_{n,l}$ is the Banach space of real valued continuous periodic functions

$$(3.2) \quad \begin{aligned} \Omega_{n,l} &= \{\omega_{n,l} = (\omega_{l'})_{l' \in \Lambda_{n,l}} \mid \omega_{l'} \in \Omega\}, \\ \Omega &= \{\omega \in C(\mathcal{I}_\beta \rightarrow \mathbb{R}) \mid \omega(0) = \omega(\beta)\}. \end{aligned}$$

The probability measure $\nu_{n,l}$ is

$$(3.3) \quad \begin{aligned} \nu_{n,l}(d\omega_{n,l}) &= \frac{1}{Z_{n,l}} \exp[-E_{n,l}(\omega_{n,l})] \chi_{n,l}(d\omega_{n,l}), \\ Z_{n,l} &= \int_{\Omega_{n,l}} \exp[-E_{n,l}(\omega_{n,l})] \chi_{n,l}(d\omega_{n,l}). \end{aligned}$$

The functions $E_{n,l} : \Omega_{n,l} \rightarrow \mathbb{R}$ are (c.f., (2.5))

$$(3.4) \quad \begin{aligned} E_{n,j}(\omega_{n,j}) &= -\frac{1}{2} \theta \varkappa^{-n(1+\delta)} \int_0^\beta \left(\sum_{l' \in \Lambda_{n,l}} \omega_{l'}(\tau) \right)^2 d\tau + \sum_{s \in \Lambda_{1,l}} E_{n-1,s}(\omega_{n-1,s}), \\ E_{0,s}(\omega_s) &= \int_0^\beta \left(\frac{a-1}{2} [\omega_s(\tau)]^2 + b [\omega_s(\tau)]^4 \right) d\tau. \end{aligned}$$

We consider $\omega_{n,l}$ as vectors $(\omega_{n-k,s})_{s \in \Lambda_{k,l}}$ with $k = 1, 2, \dots, n$ and write ω_s for $\omega_{0,s}$. The measure $\chi_{n,l}$ is

$$(3.5) \quad \chi_{n,l}(d\omega_{n,l}) = \bigotimes_{s \in \Lambda_{n,l}} \chi(d\omega_s).$$

where χ is a Gaussian measure on $\Omega_{0,s} = \Omega$. Let \mathcal{E} be the real Hilbert space $L^2(\mathcal{I}_\beta)$. Then the Banach space of continuous periodic functions Ω can be considered, up to embedding, as a subset of \mathcal{E} . The following family

$$(3.6) \quad e_q(\tau) = \begin{cases} \sqrt{\frac{2}{\beta}} \cos q\tau, & q > 0; \\ -\sqrt{\frac{2}{\beta}} \sin q\tau, & q < 0; \\ 1/\sqrt{\beta}, & q = 0. \end{cases}$$

with q varying in the set

$$(3.7) \quad \mathcal{Q} = \{q \mid q = \frac{2\pi}{\beta} \kappa, \kappa \in \mathbb{Z}\},$$

is a base of \mathcal{E} . Given $q \in \mathcal{Q}$, let P_q be the orthonormal projection on e_q . We define χ to be the Gaussian measure³ on \mathcal{E} with zero mean and with the covariance operator

$$(3.8) \quad S = \sum_{q \in \mathcal{Q}} \frac{1}{mq^2 + 1} P_q.$$

One can show (see Lemma 2.2 of [6]) that the measure χ is concentrated on Ω , i.e., $\chi(\Omega) = 1$. On the other hand, as follows from the Kuratowski theorem (see Theorem 3.9, page 21 of [22]), the Borel σ -algebras of subsets of Ω , generated by its own topology and by the topology induced from the Hilbert space \mathcal{E} , coincide. Hence, one can consider χ also as a measure on Ω . As such one, it appears in the representation (3.5).

The fluctuation operator $Q_{n,l}$, defined by (2.15) with $\lambda_n = \varkappa^{-n\delta/2}$ is a multiplication operator by the function $Q_{n,l} : \mathbb{R}^{|\Lambda_{n,l}|} \rightarrow \mathbb{R}$

$$(3.9) \quad Q_{n,l}(\xi_{n,l}) = \varkappa^{-n(1+\delta)/2} \sum_{s \in \Lambda_{n,l}} \xi_s = \varkappa^{-(1+\delta)/2} \sum_{s \in \Lambda_{1,l}} Q_{n-1,s}(\xi_{n-1,s}).$$

The representation (3.1) and the properties of the measures $\nu_{n,l}$, $\chi_{n,l}$, χ (see Lemma 2.3 and the whole section 2.2 of [6]) yield the following statement.

Proposition 3.1. *For every fixed $\beta > 0$, $\tau_1, \dots, \tau_{2k} \in \mathcal{I}_\beta$, the Matsubara functions (2.17) continuously depend on $\mathbf{m} > 0$, $a \in \mathbb{R}$ and $b > 0$.*

Proposition 3.2. *For all $n \in \mathbb{N}_0$ and $k \in \mathbb{N}$, the functions (2.17) obey the estimates*

$$(3.10) \quad 0 \leq \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) \leq \sum_{\sigma} \prod_{l=1}^k \Gamma_2^{(n)}(\tau_{\sigma(2l-1)}, \tau_{\sigma(2l)}),$$

which hold for all $\tau_1, \dots, \tau_{2k} \in \mathcal{I}_\beta$. Here the sum is taken over all possible partitions of the set $\{1, \dots, 2k\}$ onto unordered pairs.

The estimates (3.10) were proven in [6] as Theorems 6.2 (positivity) and 6.4 (Gaussian upper bound).

³For a topological space, ‘measure defined on the space’ means that the measure is defined on its Borel σ -algebra.

Since to prove our theorem we need the Matsubara functions corresponding to the operators $Q_{n,l}$ only, we may restrict our study to the measures describing distributions of $Q_{n,l}$ given by (3.9). For $n \in \mathbb{N}_0$ and a Borel subset $C \subset \Omega$, let

$$B_C = \{\omega_{n,l} \in \Omega_{n,l} \mid \varkappa^{-n(1+\delta)/2} \sum_{s \in \Lambda_{n,l}} \omega_s \in C\},$$

which is a Borel subset of $\Omega_{n,l}$. Then we set

$$\mu_n(C) = \nu_{n,l}(B_C),$$

which defines a measure on Ω . By (3.3), (3.4), the measures μ_n obey the following recursion relation

$$(3.11) \quad \mu_n(d\omega) = \frac{1}{Z_n} \exp\left(\frac{\theta}{2} \|\omega\|_{\mathcal{E}}^2\right) \mu_{n-1}^{\star \varkappa}(\varkappa^{(1+\delta)/2} d\omega),$$

$$(3.12) \quad \mu_0(d\omega) = \frac{1}{Z_0} \exp(-E_{0,s}(\omega)) \chi(d\omega),$$

where $\|\cdot\|_{\mathcal{E}}$ is the norm in the Hilbert space $\mathcal{E} = L^2(\mathcal{I}_\beta)$, the function $E_{0,s}$ is given by (3.4), Z_n , $n \in \mathbb{N}$ are normalizing constants and \star stands for convolution. For obvious reasons, we drop the labels l and s . Like the measure χ , all μ_n , $n \in \mathbb{N}_0$ can be considered either as measures on the Hilbert space \mathcal{E} concentrated on its subset Ω , or as measures on the Banach space Ω . We have

$$(3.13) \quad \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) = \int_{\Omega} \omega(\tau_1) \cdots \omega(\tau_{2k}) \mu_n(d\omega),$$

and the function (1.7) may be written in the form

$$(3.14) \quad \varphi_n^{(\delta)}(x) \stackrel{\text{def}}{=} \varphi_n(x) = \int_{\mathcal{E}} \exp((x, \omega)_{\mathcal{E}}) \mu_n(d\omega) = \int_{\Omega} \exp((x, \omega)_{\mathcal{E}}) \mu_n(d\omega), \quad x \in \mathcal{E},$$

where $(\cdot, \cdot)_{\mathcal{E}}$ is the scalar product in \mathcal{E} . Expanding its logarithm into the series (1.8) we obtain the Ursell functions (c.f., (2.17))

$$(3.15) \quad U_{2k}^{\delta, \beta, \Lambda_{n,s}}(\tau_1, \dots, \tau_{2k}) \stackrel{\text{def}}{=} U_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}), \quad k \in \mathbb{N}.$$

Correspondingly, the numbers (1.11) obtained from these functions are denoted by $\mathcal{U}_{2k}^{(n)}$. Each function $U_{2k}^{(n)}$ can be written as a polynomial of the Matsubara functions $\Gamma_{2s}^{(n)}$, $s = 1, 2, \dots, k$ and vice versa. In particular,

$$(3.16) \quad \begin{aligned} U_2^{(n)}(\tau_1, \tau_2) &= \Gamma_2^{(n)}(\tau_1, \tau_2), \\ U_4^{(n)}(\tau_1, \dots, \tau_4) &= \Gamma_4^{(n)}(\tau_1, \dots, \tau_4) - \Gamma_2^{(n)}(\tau_1, \tau_2) \Gamma_2^{(n)}(\tau_3, \tau_4) \\ &\quad - \Gamma_2^{(n)}(\tau_1, \tau_3) \Gamma_2^{(n)}(\tau_2, \tau_4) - \Gamma_2^{(n)}(\tau_1, \tau_4) \Gamma_2^{(n)}(\tau_2, \tau_3). \end{aligned}$$

In view of (2.14), the Matsubara and Ursell functions depend only on the periodic distances between τ_j , i.e., on $|\tau_i - \tau_j|_{\beta} = \min\{|\tau_i - \tau_j|, \beta - |\tau_i - \tau_j|\}$.

The proof of Theorem 2.2 is based on inequalities for the Matsubara and Ursell functions, which we obtain by means of the lattice approximation method. Its main idea is to construct sequences of probability measures, concentrated on finite dimensional subspaces of $\Omega_{n,l}$, which converge to the Euclidean measures $\nu_{n,l}$ in such a way that the integrals (3.1) are the limits of the corresponding integrals taken with such approximating measures. Then the latter integrals are being rewritten as moments of Gibbs measures of classical ferromagnetic models, for which one has

a number of useful inequalities. In such a way, these inequalities are transferred to the Matsubara and Ursell functions. A detailed description of this method is given in Section 5 of [6]. Here we provide a short explanation of its main elements. Given $N = 2L$, $L \in \mathbb{N}$, set

$$(3.17) \quad \lambda_q^{(N)} = \left\{ \mathfrak{m} \left(\frac{2N}{\beta} \right)^2 \left[\sin \left(\frac{\beta}{2N} \right) q \right]^2 + 1 \right\}^{-1},$$

and

$$(3.18) \quad S_N = \sum_{q \in \mathcal{Q}_N} \lambda_q^{(N)} P_q, \quad \mathcal{Q}_N = \{q = \frac{2\pi}{\beta} \kappa \mid \kappa = -(L-1), \dots, L\},$$

where the projectors P_q are the same as in (3.8). Now let χ_N be the Gaussian measure on \mathcal{E} with the covariance operator S_N . Let also $\chi_{n,l}^{(N)}$ be defined by (3.5) with χ_N instead of χ . By means of $\chi_{n,l}^{(N)}$, we define $\nu_{n,l}^{(N)}$ via (3.3). Then by Theorem 5.1 of [6], one has

$$(3.19) \quad \int_{\Omega_{n,l}} Q_{n,l}(\omega_{n,l}(\tau_1)) \cdots Q_{n,l}(\omega_{n,l}(\tau_{2k})) \nu_{n,l}^{(N)}(d\omega_{n,l}) \longrightarrow \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}),$$

pointwise on \mathcal{I}_β^{2k} as $N \rightarrow +\infty$. On the other hand, one can write

$$(3.20) \quad \text{LHS}(3.19) = C_{2k,N} \sum_{\ell_1, \dots, \ell_{2k}} \langle S_{\ell_1} \cdots S_{\ell_{2k}} \rangle,$$

where $C_{2k,N} > 0$ is a constant and $\langle \cdot \rangle$ stands for the expectation with respect to the local Gibbs measure on $\Xi_{n,l}^{(N)} \stackrel{\text{def}}{=} \Lambda_{n,l} \times \{1, 2, \dots, N\}$ of a ferromagnetic model with the one dimensional ϕ^4 single-spin distribution. This type of single-spin distribution is determined by our choice of the potential energy in (1.1), whereas the ferromagnetism is due to the fact that $J > 0$ (see (1.5) and due to our choice of the numbers (3.17). The sum in (3.20) is taken over the vectors $\ell_j = (\ell_j^{(1)}, \ell_j^{(2)})$, $j = 1, \dots, 2k$ as follows. Their first components run through $\Lambda_{n,l}$ and the second components are fixed at certain values from the set $\{1, \dots, N\}$, determined by the corresponding τ_j . Furthermore, the above expectations $\langle \cdot \rangle$ can be approximated by expectations with respect to the ferromagnetic Ising model (classical Ising approximation [25, 26]). Then the functions $\Gamma_{2k}^{(n)}$ and $U_{2k}^{(n)}$ obey the inequalities which the moments and semi-invariants of the ferromagnetic Ising model do obey. In particular, we have the following.

Lemma 3.3. *For all $n \in \mathbb{N}_0$ and $k \in \mathbb{N}$, the following estimates hold for all values of the arguments $\tau, \tau', \tau_1, \dots, \tau_{2k} \in \mathcal{I}_\beta$,*

$$(3.21) \quad \int_{\mathcal{I}_\beta^2} U_4^{(n)}(\tau, \tau, \tau_1, \tau_2) d\tau_1 d\tau_2 \leq \int_{\mathcal{I}_\beta^2} U_4^{(n)}(\tau, \tau', \tau_1, \tau_2) d\tau_1 d\tau_2;$$

$$(3.22) \quad (-1)^{k-1} U_{2k}^{(n)}(\tau_1, \tau_2, \dots, \tau_{2k}) \geq 0.$$

Proof. For classical models with unbounded spins and polynomial anharmonicity of the Ellis-Monroe type (for ϕ^4 -models, in particular), (3.21) was proved in [2]. For the Ising model, the sign rule (3.22) was proved in [24]. \square \square

Lemma 3.4. *For all $n, l \in \mathbb{N}_0$, the function*

$$(3.23) \quad \begin{aligned} f_n(z) &= \int_{\Omega_{n,l}} \exp\left(z \int_0^\beta Q_{n,l}(\omega_{n,l}(\tau)) d\tau\right) \nu_{n,l}(d\omega_{n,l}) \\ &= \int_{\mathcal{E}} \exp\left(z \int_0^\beta \omega(\tau) d\tau\right) \mu_n(d\omega), \end{aligned}$$

can be analytically continued to an even entire function of order less than two, possessing purely imaginary zeros.

Proof. For the function (3.23), one can construct the lattice approximation (c.f., (3.20))

$$(3.24) \quad f_n^{(N)}(z) = \left\langle \exp\left(z \sum_{\ell \in \Xi_{n,l}^{(N)}} S_\ell\right) \right\rangle,$$

which converges, as $N \rightarrow +\infty$, to $f_n(z)$ for every $z \in \mathbb{R}$. For such $f_n^{(N)}$, the property stated is known as the generalized Lee-Yang theorem [19]. The functions $f_n^{(N)}$ are ridge (crested), with the ridge being the real axis. For sequences of such functions, their pointwise convergence on the ridge implies via the Vitali theorem (see e.g., Proposition VIII.19 in [26]) the uniform convergence on compact subsets of \mathbb{C} , which by the Hurwitz theorem (see e.g., [10]) gives the desired property of f_n . \square

Set

$$(3.25) \quad \begin{aligned} \hat{u}_n(q) &= \int_0^\beta U_2^{(n)}(\tau', \tau) \cos(q\tau) d\tau, \\ &= \int_0^\beta U_2^{(n)}(0, \tau) \cos(q\tau) d\tau, \quad q \in \mathcal{Q}, \quad n \in \mathbb{N}_0. \end{aligned}$$

Then

$$(3.26) \quad U_2^{(n)}(\tau_1, \tau_2) = \frac{1}{\beta} \sum_{q \in \mathcal{Q}} \hat{u}_n(q) \cos[q(\tau_1 - \tau_2)].$$

Furthermore, we set (c.f., (1.11))

$$(3.27) \quad \mathcal{U}_{2k}^{(n)} = \int_{\mathcal{I}_\beta^{2k}} U_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) d\tau_1 \cdots d\tau_{2k}.$$

Then

$$(3.28) \quad \hat{u}_n(0) = \hat{u}_n \stackrel{\text{def}}{=} \beta^{-1} \mathcal{U}_2^{(n)} = \beta^{-1} \mathcal{U}_2^{\delta, \beta, \Lambda_n}.$$

Lemma 3.5. *For every $n \in \mathbb{N}_0$ and $q \in \mathcal{Q}$, $\hat{u}_n(q)$ is a continuous function of β , it obeys the following estimates*

$$(3.29) \quad 0 < \hat{u}_n(q) \leq \hat{u}_n;$$

$$(3.30) \quad \hat{u}_n(q) \leq \frac{\varkappa^{-n\delta}}{\mathfrak{m}q^2}, \quad q \neq 0.$$

Proof. By (3.25), (3.16), (2.17) and (2.13), one obtains

$$U_2^{(n)}(0, \tau) = \frac{1}{Z_{n,l}} \text{trace} \{ Q_{n,l} \exp[-\tau H_{n,l}] Q_{n,l} \exp[-(\beta - \tau) H_{n,l}] \}.$$

It may be shown that every $H_{n,l}$ has a pure point spectrum $\{E_p^{(n)}\}_{p \in \mathbb{N}_0}$. We denote the corresponding eigenfunctions by $\Psi_p^{(n)}$ and set

$$Q_{pp'}^{(n)} = (Q_{n,l} \Psi_p^{(n)}, \Psi_{p'}^{(n)})_{\mathcal{H}_{n,l}}.$$

Then the above representation may be rewritten

$$U_2^{(n)}(0, \tau) = \frac{1}{Z_{n,l}} \sum_{p, p' \in \mathbb{N}_0} |Q_{pp'}^{(n)}|^2 \exp \left[-\beta E_p^{(n)} + \tau (E_p^{(n)} - E_{p'}^{(n)}) \right],$$

which yields via (3.25)

$$(3.31) \quad \begin{aligned} \hat{u}_n(q) &= \frac{1}{Z_{n,l}} \sum_{p, p' \in \mathbb{N}_0} |Q_{pp'}^{(n)}|^2 \frac{E_p^{(n)} - E_{p'}^{(n)}}{q^2 + (E_p^{(n)} - E_{p'}^{(n)})^2} \\ &\quad \times \left(\exp[-\beta E_{p'}^{(n)}] - \exp[-\beta E_p^{(n)}] \right), \\ Z_{n,l} &= \sum_{p \in \mathbb{N}_0} \exp[-\beta E_p^{(n)}]. \end{aligned}$$

Both series above converge uniformly, as functions of β , on compact subsets of $(0, +\infty)$, which yields continuity and positivity. The upper bound (3.29) follows from (3.31) or from (3.25). To prove (3.30), we estimate the denominator in (3.31) from below by $q^2 \neq 0$ and obtain

$$(3.32) \quad \begin{aligned} \hat{u}_n(q) &\leq \frac{1}{q^2} \frac{1}{Z_{n,l}} \sum_{p, p'} |Q_{pp'}^{(n)}|^2 (E_p^{(n)} - E_{p'}^{(n)}) \left(\exp[-\beta E_{p'}^{(n)}] - \exp[-\beta E_p^{(n)}] \right) \\ &= \frac{1}{q^2} \frac{1}{Z_{n,l}} \text{trace} \{ [Q_{n,l}, [H_{n,l}, Q_{n,l}]] \exp(-\beta H_{n,l}) \}, \quad q \neq 0. \end{aligned}$$

By means of (2.5) and (1.2), the double commutator in (3.32) may be computed explicitly. It equals to $|\Lambda_{n,l}|^{-\delta} / \mathfrak{m}$, which yields (3.30). \square

Lemma 3.6. *The numbers $\mathcal{U}_{2k}^{(n)}$ defined by (3.27) obey the estimates (c.f., (1.16))*

$$(3.33) \quad |\mathcal{U}_{2k}^{(n)}| \leq 2^{1-k} (2k-1)! (\beta \hat{u}_n)^k, \quad k \in \mathbb{N},$$

$$(3.34) \quad |\mathcal{U}_{2k}^{(n)}| \leq \frac{(2k-1)!}{3 \cdot 2^{k-1}} (\beta \hat{u}_n)^{k-2} |\mathcal{U}_4^{(n)}|, \quad k \geq 2.$$

Proof. The function (3.23) is the same as in (1.12), hence, it possesses the representation (1.13) and $\mathcal{U}_{2k}^{(n)} = \mathcal{U}_{2k}^{\delta, \beta, \Lambda_{n,l}}$ are given by the right-hand side of (1.14). Then the estimates (3.33), (3.34) immediately follow from (1.16). \square

4. PROOF OF THE THEOREM

Set

$$(4.1) \quad X_n = - \int_{\mathcal{I}_\beta^2} U_4^{(n)}(\tau, \tau, \tau_1, \tau_2) d\tau_1 d\tau_2.$$

Then by Lemma 3.3, one has

$$(4.2) \quad 0 < \beta^{-2} \left| \mathcal{U}_4^{(n)} \right| \leq X_n, \quad \text{for all } n \in \mathbb{N}_0,$$

thus, we may control the sequence $\{\mathcal{U}_4^{(n)}\}_{n \in \mathbb{N}_0}$ by controlling $\{X_n\}_{n \in \mathbb{N}_0}$.

Lemma 4.1. *For the model (1.1) with $\delta \in (0, 1/2)$, one can choose the parameters a, b and \mathfrak{m} in such a way that there will exist $\beta_* > 0$, dependent on a, b, \mathfrak{m} , with the following properties: (a) for $\beta \leq \beta_*$, $\{X_n\}_{n \in \mathbb{N}_0} \rightarrow 0$; (b) for $\beta = \beta_*$, $\{\hat{u}_n\}_{n \in \mathbb{N}_0} \rightarrow 1$; for $\beta < \beta_*$, there exists $K(\beta) > 0$ such that for all $n \in \mathbb{N}_0$,*

$$(4.3) \quad \hat{u}_n \leq K(\beta) \varkappa^{-n\delta}.$$

The proof of this lemma will be given in the concluding section of the article. Lemmas 3.5 and 4.1 have two important corollaries.

Corollary 4.2. *For every $\beta \leq \beta_*$ and $k \in \mathbb{N}$, the sequences $\{\Gamma_{2k}^{(n)}\}_{n \in \mathbb{N}_0}$, $\{U_{2k}^{(n)}\}_{n \in \mathbb{N}_0}$ are relatively compact in the topology of uniform convergence on \mathcal{I}_β^{2k} .*

Proof. Since the Ursell functions $U_{2k}^{(n)}$ may be expressed as polynomials of $\Gamma_{2s}^{(n)}$ with $s = 1, \dots, k$ and vice versa, it is enough to prove this statement for the Matsubara functions only. By Ascoli's theorem (see e.g., [21] p. 72) we have to show that the sequence $\{\Gamma_{2k}^{(n)}\}_{n \in \mathbb{N}_0}$ is pointwise bounded and equicontinuous. By (3.30) and (3.26),

$$(4.4) \quad \Gamma_2^{(n)}(\tau, \tau') \leq \Gamma_2^{(n)}(0, 0) \leq \frac{1}{\beta} \hat{u}_n + \frac{\varkappa^{-n\delta}}{\beta \mathfrak{m}} \sum_{q \in \mathcal{Q} \setminus \{0\}} \frac{1}{q^2}.$$

For $\beta \leq \beta_*$, the sequence $\{\hat{u}_n\}_{n \in \mathbb{N}_0}$ is bounded by Lemma 4.1. Together with the Gaussian upper bound (3.10) this yields the uniform boundedness of $\Gamma_{2k}^{(n)}$ on \mathcal{I}_β^{2k} . Further, by (3.13)

$$(4.5) \quad \begin{aligned} & \Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) - \Gamma_{2k}^{(n)}(\vartheta_1, \dots, \vartheta_{2k}) \\ &= \int_{\mathcal{E}} \sum_{l=1}^{2k} \omega(\tau_1) \cdots \omega(\tau_{l-1}) [\omega(\tau_l) - \omega(\vartheta_l)] \omega(\vartheta_{l+1}) \cdots \omega(\vartheta_{2k}) \mu_n(d\omega). \end{aligned}$$

Applying here the Schwarz inequality (as to the scalar product in $L^2(\mathcal{E}, \mu_n)$ of $[\omega(\tau_l) - \omega(\vartheta_l)]$ and the rest of ω), the Gaussian upper bound (3.10) and the left-hand inequality in (4.4) one gets

$$(4.6) \quad \begin{aligned} & |\Gamma_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}) - \Gamma_{2k}^{(n)}(\vartheta_1, \dots, \vartheta_{2k})|^2 \\ & \leq \left(\Gamma_2^{(n)}(0, 0) - \Gamma_2^{(n)}(\tau, \vartheta) \right) \cdot \frac{8k^2(4k-2)!}{(2k-1)!2^{2k-1}} \left(\Gamma_2^{(n)}(0, 0) \right)^{2k-1}, \end{aligned}$$

where (τ, ϑ) is chosen amongst the pairs (τ_l, ϑ_l) , $l = 1, \dots, 2k$ to obey $|\tau - \vartheta|_\beta = \max_l |\tau_l - \vartheta_l|_\beta$. But by (3.26), (3.30),

$$\begin{aligned} \Gamma_2^{(n)}(0, 0) - \Gamma_2^{(n)}(\tau, \vartheta) &= \frac{2}{\beta} \sum_{q \in \mathcal{Q}} \hat{u}_n(q) \{\sin [(q/2)(\tau - \vartheta)]\}^2 \\ &\leq 2 \frac{\varkappa^{-n\delta}}{\beta \mathfrak{m}} \sum_{q \in \mathcal{Q} \setminus \{0\}} \frac{1}{q^2} \{\sin [(q/2)(\tau - \vartheta)]\}^2 \\ &\leq C \varkappa^{-n\delta} |\tau - \vartheta|_\beta, \end{aligned}$$

with an appropriate $C > 0$. \square

The next fact follows immediately from (3.30) and (3.7).

Corollary 4.3. *For every β ,*

$$\sum_{q \in \mathcal{Q} \setminus \{0\}} \hat{u}_n(q) \longrightarrow 0, \quad n \rightarrow +\infty.$$

Proof of Theorem 2.2. By Lemma 4.1, (3.34), and (4.1), (4.2), one obtains that for all $k \geq 2$ and $\beta \leq \beta_*$, $\{\mathcal{U}_{2k}^{(n)}\}_{n \in \mathbb{N}_0} \rightarrow 0$. Then by the sign rule (3.21), for all $k \geq 2$, the sequences $\{U_{2k}^{(n)}\}_{n \in \mathbb{N}_0}$ converge to zero for almost all $(\tau_1, \dots, \tau_{2k}) \in \mathcal{I}_\beta^{2k}$, which, by Corollary 4.2, yields their uniform convergence to zero. By (3.26) – (3.30), Corollary 4.3 and Lemma 4.1, one has for $\beta = \beta_*$,

$$(4.7) \quad U_2^{(n)}(\tau_1, \tau_2) = \frac{1}{\beta} \hat{u}_n + \frac{1}{\beta} \sum_{q \in \mathcal{Q} \setminus \{0\}} \hat{u}_n(q) \cos[q(\tau_1 - \tau_2)] \longrightarrow 1/\beta_*,$$

uniformly on \mathcal{I}_β^2 . Now one can express each $\Gamma_{2k}^{(n)}$ polynomially by $U_{2l}^{(n)}$ with $l = 1, \dots, k$ and obtain the convergence (2.18) for $\beta = \beta_*$. For $\beta < \beta_*$, we have the estimate (4.3), which yields (c.f., (4.4))

$$(4.8) \quad \Gamma_2^{\alpha, \beta, \Lambda_{n,l}}(\tau, \tau') \leq \Gamma_2^{\alpha, \beta, \Lambda_{n,l}}(0, 0) \leq \frac{\varkappa^{-n\alpha}}{\beta} \left[K(\beta) + \frac{1}{\mathfrak{m}} \sum_{q \in \mathcal{Q} \setminus \{0\}} \frac{1}{q^2} \right],$$

hence $\Gamma_2^{\alpha, \beta, \Lambda_{n,l}}(\tau, \tau') \rightarrow 0$ as $n \rightarrow +\infty$, uniformly on \mathcal{I}_β^2 . The convergence of the Matsubara functions $\Gamma_{2k}^{\alpha, \beta, \Lambda_{n,l}}$ with $k \geq 2$ follows from the Gaussian upper bound (3.10). \square

5. PROOF OF LEMMA 4.1

Set

$$(5.1) \quad \sigma(v) = \frac{\varkappa^{-\delta}}{1 - (1 - \varkappa^{-\delta})v}, \quad v \in (0, (1 - \varkappa^{-\delta})^{-1}),$$

and

$$(5.2) \quad \phi(v) = \varkappa^{2\delta-1} [\sigma(v)]^4, \quad \psi(v) = \frac{1}{2} \varkappa^{2\delta-1} (1 - \varkappa^{-\delta}) [\sigma(v)]^3.$$

Lemma 5.1. *Given $n \in \mathbb{N}$, let the condition*

$$(5.3) \quad \hat{u}_{n-1}(1 - \varkappa^{-\delta}) < 1,$$

be satisfied. Then the following inequalities hold:

$$(5.4) \quad \hat{u}_n < \sigma(\hat{u}_{n-1}) \hat{u}_{n-1};$$

$$(5.5) \quad \hat{u}_n \geq \sigma(\hat{u}_{n-1}) \hat{u}_{n-1} - \psi(\hat{u}_{n-1}) X_{n-1};$$

$$(5.6) \quad 0 < X_n \leq \phi(\hat{u}_{n-1}) X_{n-1};$$

where $\sigma(v)$, $\psi(v)$, $\phi(v)$ and X_n are defined by (5.1), (5.2) and (4.1) respectively.

Proof. For $t \in [0, \theta]$, $\theta = \varkappa^\delta - 1$, $x \in \mathcal{E}$ and $n \in \mathbb{N}$, we set (c.f., (3.14))

$$(5.7) \quad \varphi_n(x|t) = \frac{1}{Z_n} \int_{\mathcal{E}} \exp\left((x, \omega)_{\mathcal{E}} + \frac{t}{2} \|\omega\|_{\mathcal{E}}^2\right) \mu_{n-1}^{\varkappa} \left(\varkappa^{(1+\delta)/2} d\omega \right),$$

where Z_n is the same as in (3.11). Then

$$(5.8) \quad \varphi_n(x|\theta) = \varphi_n(x), \quad \varphi_n(x|0) = Z_n^{-1} \left[\varphi_{n-1} \left(\varkappa^{-(1+\delta)/2} x \right) \right]^{\varkappa}.$$

For every $t \in [0, \theta]$, the function (5.7) can be expanded in the series (1.7) with the coefficients

$$(5.9) \quad \begin{aligned} \varphi_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}|t) &= \frac{1}{Z_n} \int_{\mathcal{E}} \omega(\tau_1) \cdots \omega(\tau_{2k}) \exp\left(\frac{t}{2} \|\omega\|_{\mathcal{E}}^2\right) \\ &\times \mu_{n-1}^{\varkappa} \left(\varkappa^{(1+\delta)/2} d\omega \right), \end{aligned}$$

which, by (3.13), coincide with the corresponding Matsubara functions for $t = \theta$. For every fixed $(\tau_1, \dots, \tau_{2k}) \in \mathcal{I}_{\beta}^{2k}$, as functions of t they are differentiable at any $t \in (0, \theta)$ and continuous on $[0, \theta]$. The corresponding derivatives are obtained from (5.9)

$$(5.10) \quad \begin{aligned} \frac{\partial}{\partial t} \varphi_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}|t) &\stackrel{\text{def}}{=} \dot{\varphi}_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}|t) \\ &= \frac{1}{2} \int_0^{\beta} \varphi_{2k+2}^{(n)}(\tau_1, \dots, \tau_{2k}, \tau, \tau|t) d\tau. \end{aligned}$$

Now we write $\log \varphi_n(x|t)$ in the form of the series (1.8) and obtain the Ursell function $u_{2k}^{(n)}(\tau_1, \dots, \tau_{2k}|t)$. The derivatives of these functions with respect to t are being calculated from (5.10). In particular, this yields

$$(5.11) \quad \begin{aligned} \dot{u}_2^{(n)}(\tau_1, \tau_2|t) &= \\ &= \frac{1}{2} \int_0^{\beta} u_4^{(n)}(\tau_1, \tau_2, \tau, \tau|t) d\tau + \int_0^{\beta} u_2^{(n)}(\tau_1, \tau|t) u_2^{(n)}(\tau_2, \tau|t) d\tau; \end{aligned}$$

$$(5.12) \quad \begin{aligned} \dot{u}_4^{(n)}(\tau_1, \tau_2, \tau_3, \tau_4|t) &= \\ &= \frac{1}{2} \int_0^{\beta} u_6^{(n)}(\tau_1, \tau_2, \tau_3, \tau_4, \tau, \tau|t) d\tau + \\ &+ \int_0^{\beta} u_4^{(n)}(\tau_1, \tau_2, \tau_3, \tau|t) u_2^{(n)}(\tau_4, \tau|t) d\tau + \\ &+ \int_0^{\beta} u_4^{(n)}(\tau_1, \tau_2, \tau_4, \tau|t) u_2^{(n)}(\tau_3, \tau|t) d\tau + \\ &+ \int_0^{\beta} u_4^{(n)}(\tau_1, \tau_3, \tau_4, \tau|t) u_2^{(n)}(\tau_2, \tau|t) d\tau + \\ &+ \int_0^{\beta} u_4^{(n)}(\tau_2, \tau_3, \tau_4, \tau|t) u_2^{(n)}(\tau_1, \tau|t) d\tau. \end{aligned}$$

Then for

$$(5.13) \quad v_n(t) \stackrel{\text{def}}{=} \int_0^{\beta} u_2^{(n)}(\tau_1, \tau_2|t) d\tau_2 = \int_0^{\beta} u_2^{(n)}(0, \tau|t) d\tau,$$

we obtain the following system of equations

$$(5.14) \quad \dot{v}_n(t) = \frac{1}{2}U(t) + [v_n(t)]^2,$$

$$(5.15) \quad \begin{aligned} \dot{U}(t) &= \frac{1}{2}V(t) + 2v_n(t)U(t) + \\ &+ 2 \int_{\mathcal{I}_\beta^3} u_2^{(n)}(\tau_2, \tau_3|t)u_4^{(n)}(0, \tau_1, \tau_2, \tau_3|t)d\tau_1 d\tau_2 d\tau_3, \end{aligned}$$

subject to the initial conditions (see (5.8))

$$(5.16) \quad \begin{aligned} v_n(0) &= \varkappa^{-\delta}\hat{u}_{n-1}, \\ U(0) &= \varkappa^{-2\delta-1} \int_0^\beta u_4^{(n)}(0, \tau_1, \tau_2, \tau_2|t)d\tau_1 d\tau_2 = -\varkappa^{-2\delta-1}X_{n-1}. \end{aligned}$$

Here

$$(5.17) \quad \begin{aligned} U(t) &\stackrel{\text{def}}{=} \int_{\mathcal{I}_\beta^2} u_4^{(n)}(0, \tau_1, \tau_2, \tau_2|t)d\tau_1 d\tau_2 = \\ &= \int_{\mathcal{I}_\beta^2} u_4^{(n)}(\tau, \tau, \tau_1, \tau_2|t)d\tau_1 d\tau_2, \\ V(t) &\stackrel{\text{def}}{=} \int_{\mathcal{I}_\beta^3} u_6^{(n)}(0, \tau_1, \tau_2, \tau_2, \tau_3, \tau_3|t)d\tau_1 d\tau_2 d\tau_3. \end{aligned}$$

Along with the problem (5.14), (5.16) we consider the following one

$$(5.18) \quad \dot{y}(t) = [y(t)]^2, \quad y(0) = v_n(0) = \varkappa^{-\delta}\hat{u}_{n-1}.$$

Under the condition (5.3) it has a solution

$$(5.19) \quad y(t) = \frac{\varkappa^{-\delta}\hat{u}_{n-1}}{1 - t\varkappa^{-\delta}\hat{u}_{n-1}} = \sigma((t/\theta)\hat{u}_{n-1})\hat{u}_{n-1}, \quad t \in [0, \theta].$$

The sign rule (3.22) is valid for the above $u_{2k}^{(n)}$ for all $t \in [0, \theta]$, which yields $U(t) < 0$, $V(t) > 0$. Therefore, the solution of (5.14) will be dominated⁴ by (5.19), i.e.,

$$\hat{u}_n = v_n(\theta) < y(\theta) = \sigma(\hat{u}_{n-1})\hat{u}_{n-1},$$

that gives (5.4). Further, with the help of (3.21), (3.22) the third term on the right-hand side of (5.15) may be estimated as follows

$$\begin{aligned} &2 \int_{\mathcal{I}_\beta^2} u_2^{(n)}(\tau_2, \tau_3|t) \left(\beta^{-1} \int_{\mathcal{I}_\beta^2} u_4^{(n)}(\tau, \tau_1, \tau_2, \tau_3|t)d\tau d\tau_1 \right) d\tau_2 d\tau_3 \geq \\ &\geq 2 \left(\beta^{-1} \int_{\mathcal{I}_\beta^2} u_4^{(n)}(\tau, \tau_1, \tau_2, \tau_2|t)d\tau d\tau_1 \right) \times \\ &\times \int_{\mathcal{I}_\beta^2} u_2^{(n)}(\tau_2, \tau_3|t)d\tau_2 d\tau_3 = 2v_n(t)U(t). \end{aligned}$$

Applying this in (5.15) we arrive at (recall that $U(t) < 0$ and $V(t) > 0$)

$$(5.20) \quad \frac{\dot{U}(t)}{U(t)} \leq 4y(t) = \frac{4\varkappa^{-\delta}\hat{u}_{n-1}}{1 - t\varkappa^{-\delta}\hat{u}_{n-1}}, \quad \forall t \in [0, \theta].$$

⁴A detailed presentation of methods based on differential inequalities are given in [31].

Integrating one gets

$$(5.21) \quad U(t) \geq \frac{U(0)}{[1 - t\mathcal{X}^{-\delta}\hat{u}_{n-1}]^4}, \quad \forall t \in [0, \theta],$$

which yields in turn

$$U(\theta) = -X_n \geq -\mathcal{X}^{2\delta-1} [\sigma(\hat{u}_{n-1})]^4 X_{n-1} = -\phi(\hat{u}_{n-1})X_{n-1},$$

that gives (5.6). Now we set

$$h(t) = \frac{1}{[1 + t\mathcal{X}^{-\delta}\hat{u}_{n-1}]^2} v_n \left(\frac{t}{1 + t\mathcal{X}^{-\delta}\hat{u}_{n-1}} \right) - \frac{\mathcal{X}^{-\delta}\hat{u}_{n-1}}{1 + t\mathcal{X}^{-\delta}\hat{u}_{n-1}},$$

where $t \in [0, t_{\max}]$, $t_{\max} = \theta\mathcal{X}^\delta\sigma(\hat{u}_{n-1})$. For this function, we obtain from (5.14) the following equation

$$(5.22) \quad \dot{h}(t) = \frac{1}{2[1 + t\mathcal{X}^{-\delta}\hat{u}_{n-1}]^4} U \left(\frac{t}{1 + t\mathcal{X}^{-\delta}\hat{u}_{n-1}} \right) + [h(t)]^2,$$

subject to the boundary conditions

$$(5.23) \quad h(0) = 0, \quad h(t_{\max}) = [1 - \theta\mathcal{X}^{-\delta}\hat{u}_{n-1}]^2 [v_n(\theta) - \sigma(\hat{u}_{n-1})\hat{u}_{n-1}].$$

By means of (5.20), one may show that the first term on the right-hand side of (5.22) is a monotone increasing function of $t \in [0, t_{\max}]$, which yields

$$h(t_{\max}) - h(0) \geq t_{\max}U(0)/2.$$

Taking into account (5.23) and (5.16) one obtains from the latter

$$\begin{aligned} v_n(\theta) - \sigma(\hat{u}_{n-1})\hat{u}_{n-1} &= \hat{u}_n - \sigma(\hat{u}_{n-1})\hat{u}_{n-1} \geq \\ &\geq -\frac{1}{2}(1 - \mathcal{X}^{-\delta})[\sigma(\hat{u}_{n-1})]^3 \mathcal{X}^{2\delta-1} X_{n-1}, \end{aligned}$$

that gives (5.5). \square

Now we prove a statement, which will allow us to control the initial elements in the sequences $\{\hat{u}_n\}$, $\{X_n\}$, i.e., \hat{u}_0 and X_0 . Set

$$(5.24) \quad \eta = \eta(\beta, \mathbf{m}, a, b) = \varrho_{\beta, \Lambda_{0,l}}(\mathbf{q}_l^2) = \int_{\Omega} [\omega(0)]^2 \mu_0(d\omega).$$

From now on we suppose that $a < 0$. Set also

$$(5.25) \quad f(t) = t^{-1} (1 - e^{-t}).$$

Lemma 5.2. *The following estimates hold*

$$(5.26) \quad \frac{\beta|a|}{12b} f \left(\frac{3\beta b}{\mathbf{m}|a|} \right) \leq \hat{u}_0 \leq \min \left\{ \beta\eta; \frac{\beta|a|}{8b} \left[1 + \sqrt{1 + (16b/\beta|a|)} \right] \right\},$$

$$(5.27) \quad X_0 \leq 4!b\hat{u}_0^4 \left[f \left(\frac{3\beta b}{\mathbf{m}|a|} \right) \right]^{-1}.$$

Proof. By the equations (8.81), (8.82) of [6], we get

$$(5.28) \quad \beta\eta f \left(\frac{\beta}{4\mathbf{m}\eta} \right) \leq \hat{u}_0.$$

As in [23], we use the Bogolyubov inequality

$$\frac{\beta}{2} \varrho_{\beta, \Lambda_{0,l}} \{AA^* + A^*A\} \cdot \varrho_{\beta, \Lambda_{0,l}} \{[C^*, [H, C]]\} \geq |\varrho_{\beta, \Lambda_{0,l}} \{[C^*, A]\}|^2,$$

in which we set A to be the identity operator, $C = \mathfrak{p}_l$, $H = H_{0,l}$, and obtain

$$(5.29) \quad \eta \geq \frac{|a|}{12b}.$$

It is not difficult to show that the left-hand side of (5.28) is an increasing function of η ; hence, by (5.29) one gets the lower bound in (5.26). The upper bound $\hat{u}_0 \leq \beta\eta$, follows from the estimate (4.2) (positivity), (3.29) and the definition (5.24). One can show (see subsection 4.2 of [9] and subsection 3.2 of [8]) that the measure μ_0 is quasi-invariant with respect to the shifts $\omega \mapsto \omega + te_q$, $t \in \mathbb{R}$, $q \in \mathcal{Q}$, where e_q is given by (3.6). Its logarithmic derivative \mathfrak{b}_q in the direction e_q is

$$(5.30) \quad \mathfrak{b}_q(\omega) = -(\mathfrak{m}q^2 + a) \int_0^\beta e_q(\tau)\omega(\tau)d\tau - 4b \int_0^\beta e_q(\tau)[\omega(\tau)]^3 d\tau.$$

This derivative is used in the integration-by-parts formula

$$(5.31) \quad \int_\Omega \partial_q f(\omega) \mu_0(d\omega) = - \int_\Omega f(\omega) \mathfrak{b}_q(\omega) \mu_0(d\omega),$$

where

$$\partial_q f(\omega) \stackrel{\text{def}}{=} [(\partial/\partial t)f(\omega + te_q)]_{t=0},$$

and $f : \Omega \rightarrow \mathbb{R}$ can be taken

$$(5.32) \quad f(\omega) = \int_0^\beta e_q(\tau)\omega(\tau)d\tau.$$

We apply (5.31) with $q = 0$ to the function (5.32), also with $q = 0$, and obtain

$$(5.33) \quad 1 = -|a|\hat{u}_0 + \frac{4b}{\beta} \int_{\mathcal{I}_\beta^2} \Gamma_4^{(0)}(\tau, \tau, \tau, \tau') d\tau d\tau'.$$

By the GKS-inequality (see Theorem 6.2 in [6]),

$$\Gamma_4^{(0)}(\tau, \tau, \tau, \tau') \geq \Gamma_2^{(0)}(\tau, \tau)\Gamma_2^{(0)}(\tau, \tau'),$$

by which and by the estimate $\hat{u}_0 \leq \beta\eta$, we have in (5.33)

$$1 \geq -|a|\hat{u}_0 + 4b\eta\hat{u}_0 \geq -|a|\hat{u}_0 + 4b\beta^{-1}\hat{u}_0^2,$$

that is equivalent to the second upper bound in (5.26).

By means of the lattice approximation technique and the estimate (3.15) of [13], one gets

$$-U_4^{(0)}(\tau_1, \tau_2, \tau_3, \tau_4) \leq 4!b \int_0^\beta U_2^{(0)}(\tau_1, \tau)U_2^{(0)}(\tau_2, \tau)U_2^{(0)}(\tau_3, \tau)U_2^{(0)}(\tau_4, \tau)d\tau,$$

which yields

$$\begin{aligned} X_0 &\leq 4!b\hat{u}_0^2 \int_0^\beta [U_2^{(0)}(\tau, \tau')]^2 d\tau' \leq 4!b\hat{u}_0^3\beta\eta \\ &\leq 4!b\hat{u}_0^4 \left[f\left(\frac{\beta}{4\mathfrak{m}\eta}\right) \right]^{-1}, \end{aligned}$$

where we have used the upper bound for $\beta\eta$ obtained from (5.28). For f given by (5.25), one can show that $1/f(t)$ is an increasing function of t . Then the estimate (5.27) is obtained from the above one by means of (5.29). \square

Let us return to the functions (5.1), (5.2). Recall that we suppose $\delta \in (0, 1/2)$. Given $\epsilon \in (0, (1 - 2\delta)/4)$, we define $v(\epsilon)$ by the condition $\sigma(v(\epsilon)) = \varkappa^\epsilon$. An easy calculation yields

$$(5.34) \quad v(\epsilon) = \frac{\varkappa^\delta - \varkappa^{-\epsilon}}{\varkappa^\delta - 1} = 1 + \frac{1 - \varkappa^{-\epsilon}}{\varkappa^\delta - 1}.$$

Then

$$(5.35) \quad \phi(v) \leq \varkappa^{2\delta+4\epsilon-1} < 1, \quad \text{for } v \in [1, v(\epsilon)].$$

Furthermore, we set

$$(5.36) \quad w(\epsilon) = 2\varkappa^{1-\delta-2\epsilon} \cdot \frac{(\varkappa^\delta - \varkappa^{-\epsilon})(1 - \varkappa^{-\epsilon})}{(\varkappa^\delta - 1)^2},$$

$$(5.37) \quad w_{\max} = \sup_{\epsilon \in (0, (1-2\delta)/4)} w(\epsilon).$$

The function $\epsilon \mapsto w(\epsilon)$ is continuous, then for every $w < w_{\max}$, one finds $\varepsilon \in (0, (1 - 2\delta)/4)$ such that $w < w(\varepsilon)$. Set $\bar{v} = v(\varepsilon)$ and $\bar{w} = w(\varepsilon)$. Therefore, for this w , one has

$$(5.38) \quad -\psi(v)w + v\sigma(v) > v, \quad \text{for } v \in [1, \bar{v}].$$

Lemma 5.3. *The parameters $\mathfrak{m} > 0$, $a \in \mathbb{R}$ and $b > 0$ may be chosen in such a way that there will exist $\varepsilon \in (0, (1 - 2\delta)/4)$ and the numbers β_0^\pm , $0 < \beta_0^- < \beta_0^+ < +\infty$ with the following properties: (a) $\hat{u}_0 = 1$ for $\beta = \beta_0^-$ and $\hat{u}_0 < 1$ for $\beta < \beta_0^-$; (b) $\hat{u}_0 = \bar{v} = v(\varepsilon)$ for $\beta = \beta_0^+$ and $\hat{u}_0 < \bar{v}$ for $\beta < \beta_0^+$; (c) $X_0 < \bar{w} = w(\varepsilon)$ for all $\beta \in [\beta_0^-, \beta_0^+]$.*

Proof. Let us fix $\gamma = |a|/b$. Then by (5.26) and (5.25), one has

$$(5.39) \quad \frac{\mathfrak{m}\gamma^2}{36} \left[1 - \exp\left(-\frac{3\beta}{\mathfrak{m}\gamma}\right) \right] \leq \hat{u}_0 \leq \frac{\beta\gamma}{8} \left[1 + \sqrt{1 + \frac{16}{\beta\gamma}} \right],$$

which immediately yields $\hat{u}_0 \rightarrow 0$ as $\beta \rightarrow 0$. On the other hand, by taking $\mathfrak{m}\gamma^2 > 36\bar{v}$, one gets $\hat{u}_0 > \bar{v}$ for sufficiently large β . Since by Lemma 3.5, \hat{u}_0 depends on β continuously, this means that β_0^\pm , such that $\beta_0^- < \beta_0^+$, do exist. For fixed γ and \mathfrak{m} , the multiplier $[f(3\beta/\mathfrak{m}\gamma)]^{-1}$ in (5.27) is bounded as $\beta \in (0, \beta_0^+]$. Recall, that $\hat{u}_0 \leq \bar{v}$ for such β . Then, keeping γ fixed, we pick up b such that the right-hand side of (5.27) will be less than \bar{w} . \square

Lemma 5.4. *Let \mathfrak{I}_n , $n \in \mathbb{N}_0$, be the triple of statements $(\mathfrak{i}_n^1, \mathfrak{i}_n^2, \mathfrak{i}_n^3)$, where*

$$\begin{aligned} \mathfrak{i}_n^1 &= \{\exists \beta_n^+ \in [\beta_0^-, \beta_0^+] : \hat{u}_n = \bar{v}, \beta = \beta_n^+; \hat{u}_n < \bar{v}, \forall \beta < \beta_n^+\}; \\ \mathfrak{i}_n^2 &= \{\exists \beta_n^- \in [\beta_0^-, \beta_0^+] : \hat{u}_n = 1, \beta = \beta_n^-; \hat{u}_n < 1, \forall \beta < \beta_n^-\}; \\ \mathfrak{i}_n^3 &= \{\forall \beta \in (0, \beta_n^+) : X_n < \bar{w}\}. \end{aligned}$$

Then (i) \mathfrak{I}_0 is true; (ii) \mathfrak{I}_{n-1} implies \mathfrak{I}_n .

Proof. \mathfrak{I}_0 is true by Lemma 5.3. For $\beta = \beta_n^+$, $\sigma(\hat{u}_n) = \varkappa^\varepsilon$ and $\sigma(\hat{u}_n) < \varkappa^\varepsilon$ for $\beta < \beta_n^+$ (see (5.34), (5.35)). Set $\beta = \beta_{n-1}^+$, then (5.38), (5.36), (5.5), and \mathfrak{i}_{n-1}^3 yield

$$(5.40) \quad \begin{aligned} \hat{u}_n &\geq \varkappa^\varepsilon \bar{v} - \frac{1}{2}(1 - \varkappa^{-\delta})\varkappa^{3\varepsilon}\varkappa^{2\delta-1}X_{n-1} \\ &> \varkappa^\varepsilon \bar{v} \left[1 - \varkappa^{2(\varepsilon-1)+\delta}(\varkappa^\delta - 1) \frac{\bar{w}}{\bar{v}} \right] = \bar{v}. \end{aligned}$$

For $\beta = \beta_{n-1}^-$, the estimate (5.4) gives

$$(5.41) \quad \hat{u}_n < 1.$$

Taking into account Lemma 3.5 (continuity) and the estimates (5.40), (5.41), one concludes that there exists at least one value $\tilde{\beta}_n^+ \in (\beta_{n-1}^-, \beta_{n-1}^+)$ such that $\hat{u}_n = \bar{v}$. Then we put $\beta_n^+ = \min \tilde{\beta}_n^+$. The mentioned continuity of \hat{u}_n yields also $\hat{u}_n < \bar{v}$ for $\beta < \beta_n^+$. Thus i_n^1 is true. The existence of $\beta_n^- \in [\beta_{n-1}^-, \beta_{n-1}^+)$ can be proven in the same way. For $\beta < \beta_n^+ < \beta_{n-1}^+$, we have $\sigma(\hat{u}_{n-1}) < \varkappa^\varepsilon$, which yields

$$(5.42) \quad X_n < \varkappa^{2\delta-1} \varkappa^{4\varepsilon} X_{n-1} \leq X_{n-1} < \bar{w},$$

hence, i_n^3 is true as well. The proof is concluded by remarking that

$$(5.43) \quad [\beta_n^-, \beta_n^+] \subset [\beta_{n-1}^-, \beta_{n-1}^+] \subset [\beta_0^-, \beta_0^+].$$

□

□

Lemma 5.5. *There exists $\beta_* \in [\beta_0^-, \beta_0^+]$ such that, for $\beta = \beta_*$, the following estimates hold for all $n \in \mathbb{N}_0$:*

$$(5.44) \quad 1 < \hat{u}_n < \bar{v}.$$

For $\beta < \beta_*$, the above upper estimate, as well as the estimate (4.3), hold.

Proof. Consider the set $\Delta_n \stackrel{\text{def}}{=} \{\beta \in (0, \beta_n^+) \mid 1 < \hat{u}_n < \bar{v}\}$. Just above we have shown that it is nonempty and $\Delta_n \subseteq (\beta_n^-, \beta_n^+)$. Let us prove that $\Delta_n \subseteq \Delta_{n-1}$. Suppose there exists some $\beta \in \Delta_n$, which does not belong to Δ_{n-1} . For this β , either $\hat{u}_{n-1} \leq 1$ or $\hat{u}_{n-1} \geq \bar{v}$. Hence, either $\hat{u}_n < 1$ or $\hat{u}_n > \bar{v}$ (it can be proven as above), which is in conflict with the assumption $\beta \in \Delta_n$. Now let D_n be the closure of Δ_n , then one has

$$(5.45) \quad D_n = \{\beta \in [\beta_n^-, \beta_n^+] \mid 1 \leq \hat{u}_n \leq v(\delta)\},$$

which is a nonempty closed set. Furthermore, $D_n \subseteq D_{n-1} \subseteq \dots \subseteq [\beta_0^-, \beta_0^+]$. Set $D_* = \bigcap_n D_n$, then $D_* \subset [\beta_0^-, \beta_0^+]$ is also nonempty and closed. Now let us show that, for every $\beta \in D_*$, the sharp bounds in (5.44) hold for all $n \in \mathbb{N}$. Suppose $\hat{u}_n = \bar{v}$ for some $n \in \mathbb{N}$. Then (5.40) yields $\hat{u}_m > \bar{v}$ for all $m > n$, which means that this β does not belong to all D_m , and hence to D_* . Similarly one proves the lower bound by means of (5.4). On the other hand, by means of the above arguments, one can conclude that $\beta \in D_*$ if the inequalities (5.44) hold for all $n \in \mathbb{N}_0$ at this β . Set $\beta_* = \min D_*$. Then (5.44) hold for $\beta = \beta_*$. Let us prove (4.3). Take $\beta < \beta_*$. If $\hat{u}_n > 1$ for all $n \in \mathbb{N}$, then either (5.44) holds or there exists such $n_0 \in \mathbb{N}$ that $\hat{u}_{n_0} \geq \bar{v}$. Therefore, either $\beta \in D_*$ or $\beta > \inf \beta_n^+$. Both these cases contradict the assumption $\beta < \beta_*$. Thus, there exists $n_0 \in \mathbb{N}$ such that $\hat{u}_{n_0-1} \leq 1$ and hence $\hat{u}_n < 1$ for all $n \geq n_0$. In what follows, the definition (5.1) and the estimate (5.4) imply that the sequences $\{\hat{u}_n\}_{n \geq n_0}$ and $\{\sigma(\hat{u}_n)\}_{n \geq n_0}$ are strictly decreasing. Then for all $n > n_0$, one has (see (5.4))

$$\begin{aligned} \hat{u}_n &< \sigma(\hat{u}_{n-1})\hat{u}_{n-1} < \dots \\ &< \sigma(\hat{u}_{n-1})\sigma(\hat{u}_{n-2}) \dots \sigma(\hat{u}_{n_0})\hat{u}_{n_0} < [\sigma(\hat{u}_{n_0})]^{n-n_0}. \end{aligned}$$

Since $\sigma(\hat{u}_{n_0}) < 1$, one gets $\sum_{n=0}^{\infty} \hat{u}_n < \infty$. Thus,

$$\prod_{n=1}^{\infty} [1 - (1 - \varkappa^{-\delta})\hat{u}_{n-1}]^{-1} \stackrel{\text{def}}{=} K_0 < \infty.$$

Finally, we apply (5.4) once again and obtain

$$\begin{aligned}\hat{u}_n &< \varkappa^{-n\delta} [1 - (1 - \varkappa^{-\delta})\hat{u}_{n-1}]^{-1} \dots [1 - (1 - \varkappa^{-\delta})\hat{u}_0]^{-1} \hat{u}_0 \\ &< \varkappa^{-n\delta} K_0 \bar{v} \stackrel{\text{def}}{=} K(\beta) \varkappa^{-n\delta}.\end{aligned}$$

□

□

Proof of Lemma 4.1. The existence of β_* has been proven in Lemma 5.5. Consider the case $\beta = \beta_*$ where the estimates (5.44) hold. First we show that $X_n \rightarrow 0$. Making use of (5.6) we obtain

$$0 < X_n \leq \varkappa^{2\delta-1} [\sigma(\hat{u}_{n-1})]^4 X_{n-1} < X_{n-1} < X_{n-2} < \dots < \bar{v}.$$

Therefore, the sequence $\{X_n\}$ is strictly decreasing and bounded, hence, it converges and its limit, say X_* , obeys the condition $X_* < X_0 < \bar{v}$. Assume that $X_* > 0$. Then (5.6) yields $\sigma(\hat{u}_n) \rightarrow \varkappa^\varepsilon$ hence $\hat{u}_n \rightarrow \hat{u}_\infty \geq \bar{v}$. Passing to the limit $n \rightarrow \infty$ in (5.5) one obtains $X_* \geq \bar{v}$ which contradicts the above condition. Thus $X_* = 0$. To show $\hat{u}_n \rightarrow 1$ we set

$$(5.46) \quad \Xi_n = -\frac{1}{2}(1 - \varkappa^{-\delta}) [\sigma(\hat{u}_{n-1})]^3 \varkappa^{2\delta-1} X_{n-1}.$$

Combining (5.4) and (5.5) we obtain

$$(5.47) \quad 0 \geq \hat{u}_n - \sigma(\hat{u}_{n-1})\hat{u}_{n-1} \geq \Xi_n \rightarrow 0.$$

For $\beta = \beta_*$, we have $\{\hat{u}_n\} \subset [1, \bar{v}]$ in view of Lemma 5.5. By (5.47) all its accumulation points in $\in [1, \bar{v}]$ ought to solve the equation

$$u - \sigma(u)u = 0.$$

There is only one such point: $u_* = 1$, which hence is the limit of the whole sequence $\{\hat{u}_n\}$. For $\beta < \beta_*$, the estimate (4.3) has been already proven in Lemma 5.5. This yields $\sigma(\hat{u}_n) \rightarrow \varkappa^{-\delta}$, which implies $X_n \rightarrow 0$ if (5.6) is taken into account.

□

REFERENCES

- [1] Albeverio, S., Høegh-Krohn, R.: Homogeneous random fields and quantum statistical mechanics. *J. Funct. Anal.* **19**, 242-272 (1975)
- [2] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu.: Quantum hierarchical model. *Methods of Funct. Anal. Topology.* **2**, 1-35 (1996)
- [3] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu.: Critical point properties of a quantum hierarchical model. *Lett. Math. Phys.* **40**, 287-291 (1997)
- [4] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu.: Absence of critical points for a class of quantum hierarchical models. *Commun. Math. Phys.* **187**, 1-18 (1997)
- [5] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu.: Classical limits of Euclidean Gibbs states for quantum lattice models. *Lett. Math. Phys.* **48**, 221-233 (1999)
- [6] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu., Röckner, M.: Euclidean Gibbs states of quantum lattice systems. *Rev. Math. Phys.* **14**, 1335-1401 (2002)
- [7] Albeverio, S., Kondratiev, Yu., Kozitsky, Yu., Röckner, M.: Quantum stabilization in anharmonic crystals. *Phys. Rev. Lett.* **90**, No 17, 170603-1-4 (2003)
- [8] Albeverio, S., Kondratiev, Yu., Pasurek, T., Röckner, M.: Euclidean Gibbs measures on loop lattices: existence and a priori estimates. *Annals of Probab.* (to appear)
- [9] Albeverio, S., Kondratiev, Yu., Röckner, M.: Ergodicity for the stochastic dynamics of quasi-invariant measures with applications to Gibbs states. *J. Funct. Anal.* **149**, 415-469 (1997)
- [10] Berenstein, C.A., Gay, R.: *Complex Variables. An Introduction.* New York Berlin Heidelberg, Springer-Verlag 1991

- [11] Bleher, P.M.: Construction of non-Gaussian self-similar random fields with hierarchical structure. *Commun. Math. Phys.* **84**, 557-578 (1982)
- [12] Bleher, P.M., Major, P.: Critical phenomena and universal exponents in statistical physics. On Dyson's hierarchical model. *Annals of Probab.* **15**, 431-477 (1987)
- [13] Brydges D.C., Fröhlich J., Sokal A.D.: The random-walk representation of classical spin systems and correlation inequalities. II. The skeleton inequalities. *Commun. Math. Phys.* **91**, 117-139 (1983)
- [14] Freericks, J.K, Jarrel, M., Mahan, G.D.: The anharmonic electron-phonon problem. *Phys. Rev. Lett.* **77**, 4588-4591 (1996)
- [15] Høegh-Krohn, R.: Relativistic quantum statistical mechanics in two-dimensional space-time. *Commun. Math. Phys.* **38**, 195-224 (1974)
- [16] Kozitsky, Yu.: Hierarchical model of a ferromagnet. Self-similar block-spin distributions and the Lee-Yang theorem. *Rep. Math. Phys.* **26**, 429- 445 (1988)
- [17] Kozitsky, Yu.: Hierarchical ferromagnetic vector spin model possessing the Lee-Yang property. Thermodynamic limit at the critical point and above. *J. Stat. Phys.* **87**, 799-820 (1997)
- [18] Kozitsky, Yu., Wolowski L.: A nonlinear dynamical system on the set of Laguerre entire functions. *Nonlinear Anal. Ser. A: Theory, Methods.* **49**, 61-86 (2002)
- [19] Lieb, E.H., Sokal, A.D.: A general Lee-Yang theorem for one-component and multicomponent ferromagnets. *Commun. Math. Phys.* **80**, 153-179 (1981)
- [20] Moreira, C.H.C., Schor, R.C.: On the critical behavior of Dyson's quantum hierarchical models. *Commun. Math. Phys.* **160**, 275-294 (1994)
- [21] Mujica, J.: *Complex Analysis in Banach Spaces*. Amsterdam - New York - Oxford: North Holland 1986
- [22] Parthasarathy, K.R.: *Probability Measures on Metric Spaces*. New York - London: Academic Press 1967
- [23] Pastur, L.A., Khoruzhenko B.A.: Phase transitions in quantum models of rotators and ferroelectrics. *Teoret. Mat. Fiz.* **73**, 111-124 (1987)
- [24] Shlosman, S.B.: Signs of the Ising model Ursell functions. *Commun. Math. Phys.* **102**, 679-686 (1986)
- [25] Simon, B., Griffiths, R.B.: The ϕ_2^4 field theory as a classical Ising model. *Commun. Math. Phys.* **33**, 145-164 (1973)
- [26] Simon, B.: *The $P(\phi)_2$ Euclidean (Quantum) Field Theory*. Princeton, New Jersey: Princeton University Press 1974
- [27] Sinai, Ya. G.: Self-similar probability distributions. *Teor. Veroyatn. Ee Primen.* **21**, 63-80 (1976)
- [28] Sinai, Ya. G.: *Theory of Phase Transitions: Rigorous Results*. Oxford - New York: Pergamon Press 1982
- [29] Stamenković, S.: Unified model description of order-disorder and structural phase transitions. *Condensed Matter Physics (Lviv)* **1**(14), 257-309 (1998)
- [30] Stasyuk, I.V.: Local anharmonic effects in high- T_c superconductors. Pseudospin-electron model. *Condensed Matter Physics (Lviv)* **2**(19), 435-446 (1999)
- [31] Walter, W.: *Differential and Integral Inequalities*. Berlin-Heidelberg-New York: Springer-Verlag, 1970

ABTEILUNG FÜR STOCHASTIK, UNIVERSITÄT BONN, D 53115 BONN, GERMANY, BiBoS RESEARCH CENTRE, BIELEFELD, GERMANY, CERFIM, LOCARNO AND ACADEMIA DI ARCHITETTURA,, UNIVERSITÀ DELLA SVIZZERA ITALIANA, MENDRISIO, SWITZERLAND, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TRENTO, TRENTO, ITALY

E-mail address: albeverio@uni-bonn.de

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, D 33615 BIELEFELD, GERMANY, BiBoS RESEARCH CENTRE, BIELEFELD, GERMANY, INSTITUTE OF MATHEMATICS, KIEV, UKRAINE

E-mail address: kondrat@mathematik.uni-bielefeld.de

INSTYTUT MATEMATYKI, UNIwersytet Marii Curie-Skłodowskiej, PL 20-031 LUBLIN, POLAND

E-mail address: akozak@golem.umcs.lublin.pl

INSTYTUT MATEMATYKI, UNIwersytet Marii Curie-Skłodowskiej, PL 20-031 LUBLIN, POLAND

E-mail address: jkozi@golem.umcs.lublin.pl