

# INTEGRAL RATIOS OF FACTORIALS AND ALGEBRAIC HYPERGEOMETRIC FUNCTIONS

FERNANDO RODRIGUEZ-VILLEGAS

Chebychev in his work on the distribution of primes numbers used the following fact

$$u_n := \frac{(30n)!n!}{(15n)!(10n)!(6n)!} \in \mathbb{Z}, \quad n = 0, 1, 2, \dots$$

This is not immediately obvious (for example, this ratio of factorials is not a product of multinomial coefficients) but it is not hard to prove. The only proof I know proceeds by checking that the valuations  $v_p(u_n)$  are non-negative for every prime  $p$ ; an interpretation of  $u_n$  as counting natural objects or being dimensions of natural vector spaces is far from clear.

As it turns out, the generating function

$$u := \sum_{\nu \geq 1} u_n \lambda^n$$

is algebraic over  $\mathbb{Q}(\lambda)$ ; i.e. there is a polynomial  $F \in \mathbb{Z}[x, y]$  such that

$$F(\lambda, u(\lambda)) = 0.$$

However, we are not likely to see this polynomial explicitly any time soon as its degree is 483,840 (!)

What is the connection between  $u_n$  being an integer for all  $n$  and  $u$  being algebraic? Consider the more general situation

$$u_n := \prod_{\nu \geq 1} (\nu n)!^{\gamma_\nu},$$

where the sequence  $\gamma = (\gamma_\nu)$  for  $\nu \in \mathbb{N}$  consists of integers which are zero except for finitely many.

We assume throughout that  $\gamma$  is *regular*, i.e.,

$$\sum_{\nu \geq 1} \nu \gamma_\nu = 0,$$

which, by Stirling's formula, is equivalent to the generating series  $u := \sum_{\nu \geq 1} u_n \lambda^n$  having finite non-zero radius of convergence. We define the *dimension* of  $\gamma$  to be

$$d := - \sum_{\nu \geq 1} \gamma_\nu.$$

To abbreviate, we will say that  $\gamma$  is *integral* if  $u_n \in \mathbb{Z}$  for every  $n = 0, 1, 2, \dots$

We can now state the main theorem of the talk.

**Theorem 1.** *Let  $\gamma \neq 0$  be regular; then  $u$  is algebraic if and only if  $\gamma$  is integral and  $d = 1$ .*

One direction is fairly straightforward. If  $u$  is algebraic, by a theorem of Eisenstein, there exists an  $N \in \mathbb{N}$  such that  $N^n u_n \in \mathbb{N}$  for all  $n \in \mathbb{N}$ . It is not hard to see that in our case if such an  $N$  exists then it must equal 1. To see that  $d = 1$  we need to introduce the *monodromy representation*.

The power series  $u$  satisfies a linear differential equation  $Lu = 0$ . After possibly scaling  $\lambda$  this equation has singularities only at  $0, 1$  and  $\infty$ . Indeed,  $u$  is a hypergeometric series. Moreover, these singularities are regular singularities precisely because we assumed  $\gamma$  to be regular.

If we let  $V$  be the space of local solutions to  $Lu = 0$  at some base point not  $0, 1$  or  $\infty$  then analytic continuation gives a representation

$$\rho : \pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \longrightarrow GL(V).$$

We let the *monodromy group*  $\Gamma$  be the image of  $\rho$  and let  $B, A, \sigma$  be the monodromies around  $0, \infty, 1$ , respectively, with orientations chosen so that  $A = B\sigma$ . The main use of the monodromy group for us is the fact that  $u$  is algebraic if and only if  $\Gamma$  is finite.

As it happens the multiplicity of the eigenvalue 1 for  $B$  is  $d$  and it is also true that the corresponding Jordan block of  $B$  is of size  $d$ . Hence,  $\Gamma$  is not finite if  $d > 1$ .

To prove the converse we appeal to the work of Beukers and Heckman [1] who extended Schwartz work and described all algebraic hypergeometric functions. Let  $p$  and  $q$  be the characteristic polynomials of  $A$  and  $B$  respectively. In our situation  $p$  and  $q$  are relatively prime polynomials in  $\mathbb{Z}[x]$  (which are products of cyclotomic polynomials). Their work tells us that  $\Gamma$  is finite if and only if the roots of  $p$  and  $q$  interlace in the unit circle.

The key step in the proof of this beautiful fact is to determine when  $\Gamma$  fixes a non-trivial positive definite Hermitian form  $H$  on  $V$  (which guarantees that  $\Gamma$  is compact). I explained in my talk how  $H$  can be defined using a variant of a construction going back to Bezout. Consider the two variable polynomial

$$\frac{p(x)q(y) - p(y)q(x)}{x - y} = \sum_{i,j} B_{i,j} x^i y^j$$

and define the *Bezoutian* of  $p$  and  $q$  as

$$\text{Bez}(p, q) = (B_{i,j}).$$

We need two facts about this matrix. First, the determinant of  $\text{Bez}(p, q)$  equals the resultant of  $p$  and  $q$  (in passing I should mention that this is a useful fact computationally since the matrix is of smaller size than the usual Sylvester matrix). Second, note that  $\text{Bez}(p, q)$  is symmetric. Hence it carries more information than just its determinant as it defines a quadratic form  $H$ . It is a classical fact (due to Hermite and Hurwitz) that the signature of  $H$  has a topological interpretation.

Consider the continuous map  $\mathbb{P}^1(\mathbb{R}) \rightarrow \mathbb{P}^1(\mathbb{R})$  given by the rational function  $p/q$ . Since  $\mathbb{P}^1(\mathbb{R})$  is topologically a circle we have  $H^1(\mathbb{P}^1(\mathbb{R}), \mathbb{Z}) \simeq \mathbb{Z}$  and the induced map  $H^1(\mathbb{P}^1(\mathbb{R}), \mathbb{Z}) \rightarrow H^1(\mathbb{P}^1(\mathbb{R}), \mathbb{Z})$  is multiplication by some integer  $s$ , which is none other than the signature of  $H$ . In particular,  $H$  is definite if and only if the roots of  $p$  and  $q$  interlace on  $\mathbb{R}$ . A twisted form of this construction and analogous signature result can be applied to the hypergeometric situation; in this way we recover the facts about the Hermitian form fixed by  $\Gamma$  proved by Beukers and Heckman.

Finally, to make the connection with the integrality of  $\gamma$  we define the *Landau function*

$$\mathcal{L}(x) := - \sum_{\nu \geq 1} \gamma_\nu \{\nu x\}, \quad x \in \mathbb{R}$$

where  $\{x\}$  denotes fractional part. It is simple to verify that

$$v_p(u_n) = \sum_{k \geq 1} \mathcal{L}\left(\frac{n}{p^k}\right).$$

Landau [2] proved a nice criterion for integrality:  $\gamma$  is integral if and only if  $\mathcal{L}(x) \geq 0$  for all  $x \in \mathbb{R}$ .

Write

$$p(t) = \prod_{j=1}^r (t - e^{2\pi i \alpha_j}), \quad q(t) = \prod_{j=1}^r (t - e^{2\pi i \beta_j}),$$

where  $r = \dim V$  and  $0 \leq \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_r < 1$  and  $0 \leq \beta_1 \leq \beta_2 \leq \dots \leq \beta_r < 1$  are rational.

The function  $\mathcal{L}$  satisfies a number of simple properties: it is locally constant (by regularity), periodic modulo 1, right continuous with discontinuity points exactly at  $x \equiv \alpha_j \pmod{1}$  or  $x \equiv \beta_j \pmod{1}$  for some  $j = 1, \dots, r$  and takes only integer values. More precisely,

$$\mathcal{L}(x) = \#\{j \mid \alpha_j \leq x\} - \#\{j \mid 0 < \beta_j \leq x\}.$$

Away from the discontinuity points of  $\mathcal{L}$  we have

$$\mathcal{L}(-x) = d - \mathcal{L}(x).$$

In particular,  $\mathcal{L}(x) \geq 0$  if and only if  $\mathcal{L}(x) \leq d$ .

It is now easy to verify that if  $d = 1$  and  $\mathcal{L}(x) \geq 0$  then the roots of  $p$  and  $q$  must necessarily interlace on the unit circle finishing the proof. (Some further elaboration would also yield the other implication in the theorem independently of our previous argument.)

As a final note, let me mention that the examples in the theorem are a case of the ADE phenomenon; up to the obvious scaling  $n \mapsto dn$  for some  $d \in \mathbb{N}$ , they come in two infinite families  $A$  and  $D$ , which are easy to describe, and some sporadic ones (10 of type  $E_6$ , 10 of type  $E_7$  and 30 of type  $E_8$ ).

#### REFERENCES

- [1] F. Beukers and G. Heckman *Monodromy for the hypergeometric function  ${}_nF_{n-1}$* , Invent. Math. **95** (1989), 325–354.
- [2] E. Landau *Sur les conditions de divisibilité d'un produit de factorielles par un autre*. Collected works, I, p. 116, Thales-Verlag, Essen, 1985.

DEPARTMENT OF MATHEMATICS UNIVERSITY OF TEXAS AT AUSTIN, TX 78712  
*E-mail address:* villegasmath.utexas.edu